

METHODS FOR THE PRODUCTION OF STABLY TRANSFORMED, FERTILE  
GRAMINEAE EMPLOYING AGROBACTERIUM-MEDIATED TRANSFORMATION OF  
ISOLATED GRAMINEAE ZYGOTES

5 BACKGROUND OF THE INVENTION

Field of the Invention

10 The present invention relates to methods for the incorporation of DNA into the genome of a Gramineae plant, preferably a wheat plant, by means of Agrobacterium-mediated transformation. In other aspects, the invention relates to the production of stably transformed and fertile Gramineae plants, preferably wheat plants, gametes and offspring from these plants.

15 Description of the Related Art

During the past decade, it has become possible to transfer genes from a wide range of organisms to crop plants by recombinant DNA technology. This advance has provided enormous opportunities to improve plant resistance to pests, diseases and herbicides, and to modify biosynthetic processes to change the quality of plant products. However, the availability of an efficient transformation method to introduce foreign DNA has been a substantial barrier for most monocot species, including maize, rice, oat, barley, and particularly wheat.

25 There have been many methods attempted for the transformation of monocotyledonous plants, wherein "biolistics" is the most widely used transformation method for monocotyledons. In the "biolistics" (microprojectile-mediated DNA delivery) method microprojectile particles are coated with DNA and accelerated by a mechanical device to a speed high enough to penetrate the plant cell wall and nucleus (WO 91/02071).  
30 The foreign DNA gets incorporated into the host DNA and results in a transformed cell. There are many variations on the "biolistics" method (Sanford, 1990; Fromm et al., 1990; Christou et al., 1988; Sautter et al., 1991). The method has been used to produce stably transformed monocotyledonous plants including rice, maize, wheat, barley, and oats (Christou et al., 1991; Gordon-Kamm et al., 1990; Vasil et al., 1992, 1993;  
35 Wan et al., 1994; Sommers et al., 1992). However, even with the more recent improvements in transformation methods using immature embryos as target tissues, it still requires 4 to 6 months to recover transgenic plants (Weeks et al., 1993; Vasil et al., 1992; 1993; Becker et al., 1994, Rasco-Gaunt et al. 2001). The frequency of transgenic plant formation by these methods is variable ranging from less than one to more than  
40 17 events from 100 bombarded embryos (Rasco-Gaunt et al. 2001). Microprojectile-mediated DNA delivery brings about a number of problems such as frequent fragmentation of the DNA-sequence prior to its integration, random integration in transcribed as well as non-transcribed chromosomal regions, predominantly multiple insertion of the sequence to be transferred, complex integration patterns, integration of backbone  
45 sequences including selectable marker genes at the same locus. Moreover, microprojectile-mediated plant transformation is generally based upon genotype-dependent cell culture methods which often require a secondary transfer of the transgene into the background of elite breeding material via long-lasting back-crossing.

Protoplast based methods have been used mostly in rice, where DNA is delivered to the protoplasts through liposomes, PEG, or electroporation (Shimamoto 1989; Datta 1990b). Protoplasts may be isolated from various tissues but require in general the use of cell wall-degrading enzymes. It is considered likely that the use of cell wall-degrading enzymes can inhibit the subsequent regeneration process. Furthermore, most protoplast based methods require the establishment of long-term embryogenic suspension cultures. Some regenerants from protoplasts are infertile and phenotypically abnormal due to somaclonal variation during the long-term suspension culture (Davey 1991; Rhodes 1988). Transformation by electroporation involves the application of short, high-voltage electric fields to create "pores" in the cell membrane through which DNA is taken-up. This method has been used to produce stably transformed monocotyledonous plants (Paszkowski 1984; Shillito 1985; Fromm 1986) especially from rice (Shimamoto 1989; Datta 1990b; Hayakawa 1992).

A number of other methods have been reported for the transformation of monocotyledonous plants including, for example, the "pollen tube method" (WO 93/18168; Luo 1988), macro-injection of DNA into floral tillers (Du 1989; De la Pena 1987), injection of *Agrobacterium* into developing caryopses (WO 00/63398), and tissue incubation of seeds in DNA solutions (Töpfer et al. 1989). Direct injection of exogenous DNA into the fertilized plant ovule at the onset of embryogenesis was disclosed in WO 94/00583.

While widely useful in dicotyledonous plants, *Agrobacterium*-mediated gene transfer has long been disappointing when adapted to use in monocots. There are several reports in the literature claiming *Agrobacterium* transformation of monocotyledons (e.g., discussed WO 94/00977). These are specifically the methods of Gould 1991; Mooney 1991; and Raineri 1990, which claim *Agrobacterium* transformation of maize, rice and wheat. There is some evidence of gene transfer in these methods but they lack convincing evidence for transfer efficiency, reproducibility, and confirmation of gene transfer (Potrykus, 1990), and lack of transfer to the progeny when plants are produced. In the work of Gould where evidence of transformed plants was presented there was no Mendelian inheritance of the genes. Attempts by Hiei et al. (1994) suggested that transgenic rice plants could be obtained following *Agrobacterium*-mediated transformation, but the particular bacterial strains used and the choice of bacterial vectors were critical for successfully obtaining transgenics. A paper by Ishida et al. (1996) indicated that high-efficiency transformation of maize was possible by co-culture of immature embryos with *A. tumefaciens*. In both reports on rice and maize transformation, a super-binary vector pTOK233 containing additional copies of the *virB*, *virC* and *virG* genes was used to achieve high-efficiency transformation. A report by Saito et al. (WO 95/06722) disclosed a method of transforming monocotyledons using scutellum of immature embryos with *A. tumefaciens*.

WO 97/48814 disclosed a process for producing stably transformed fertile wheat and a system of transforming wheat via *Agrobacterium* based on freshly isolated or pre-cultured immature embryos, embryogenic callus and suspension cells. Also disclosed are methods for recovering transgenic plants after transformation within a short period of time, provided the explants are regenerable at the time of transformation. All methods disclosed therein use at some stage dedifferentiation of the target tissue. Selection is performed at callus stage to enable preferential development of transformed tissue eventually resulting in regeneration of homogeneously transformed plants. The number

of transgenic events obtained from the various starting materials is relatively low and generally below the 1% transformation efficacy range. This makes any effort to obtain transgenic wheat plants a laborious, cost-intensive approach.

- 5 The above mentioned plant transformation systems involve callus-based selection protocols and extensive in vitro culturing. These methodologies are time-consuming and increase the likelihood that somaclonal variants will arise that exhibit undesirable agro-  
10 nomic characteristics. Furthermore, these methods are extremely genotype-depending and need extensive adoption of the cell culture conditions for each plant species (and often even each line). Use of developmentally organized explants (such as immature embryos) as targets for transformation circumvent time-consuming tissue culture steps but increases the likelihood that chimeric plants are produced.

- 15 Targeting gametes, zygotes or early stage embryos in embryo sacs for transformation is a potential solution to these problems. WO 98/01576 discloses a method of plant (especially maize) transformation in which a foreign gene might be introduced into the zygote situated in its natural cellular context (the embryo sac). The intact embryo sac is claimed to be of essential importance for the subsequent recovery of transgenic plants. While Agrobacterium-mediated transformation of those in situ maize zygotes is hypo-  
20 thetically described, it seems unlikely that this approach will result in stably transformed plants, since Agrobacteria have only severely restricted access to the zygote within the embryo sac.

- 25 The differentiation of the zygote into an embryo capable of germination is a key process in the life cycle of plants. Investigations on embryogenesis in higher plants have been performed by either descriptive examination of morphology (reviewed by Johri et al. 1992), genetic analysis of developmental mutants (reviewed by Meinke 1993 and by Jürgens et al. 1994), or molecular analysis of gene expression (Thomas 1993). Although the production of zygotes by plants is well-understood, reproducible methods  
30 for in vitro manipulation and transformation of these cells are needed. In most plant species zygotes are comprised in embryo sacs which are deeply enclosed in the sporophytic tissues and therefore the zygotes are difficult to manipulate. Pollination involves the transfer of pollen from anther to stigma. Pollen germinates on the stigma and a pollen tube grows through the style and enters the tip of the ovule through the  
35 micropyle. Two male gametes or sperm cells are formed by division of the generative cell of the pollen grain. The sperm cells move through the pollen tube and are emptied into the embryo sac. Within each ovule there is a megasporocyte that undergoes meiosis to generate four haploid megaspores. Three of the megaspores usually disintegrate while the fourth continues to undergo divisions and forms an embryo sac which origi-  
40 nally contains six haploid cells as well as the diploid central cell. The egg and two synergids are found near the micropyle while three antipodals are found at the chalazal end of the embryo sac. One sperm cell fuses with the egg cell to form the zygote which eventually develops into the embryo and a new plant. The second sperm fuses with the central cell to form the triploid primary endosperm cell which constitutes the origine of  
45 the endosperm.

Methods for the isolation of living embryo sacs have been developed for plant species by using either micro-dissection or enzymatic digestion (Allington, 1985; Theunis 1991; Wu 1993). In vitro manipulation of fertilized embryo sacs frequently results in low viabil-

ity or production of abnormal embryos (Leduc 1995). To date, only a few reports have been issued on embryonic development and regeneration from higher-plant zygotes free from maternal tissue. Kranz and coworkers reported microcallus development (Kranz et al. 1991; Kranz and Lörz 1994), embryogenesis and plant regeneration (Kranz and Lörz 1993) from cultivated maize zygotes which had been produced by in-vitro fertilization using isolated sperm and egg cells. Formation of embryo-like structures followed by plant regeneration was also reproducibly obtained by cultivating isolated barley (Holm et al. 1994) and maize zygotes (Leduc et al. 1996) after fertilization in planta. Furthermore, methods for isolation of wheat zygotes and their regeneration into wheat plants are described (Kumlehn et al. 1997; Kumlehn et al. 1998).

WO 94/01999 is describing a method of regeneration of plants from fertilized, isolated egg protoplasts. Furthermore, a method of transformation of fertilized egg cells and regeneration of transgenic plants therefrom is described. It is explicitly stated, that grasses (Gramineae like e.g., barley or wheat) are not susceptible to Agrobacterium-mediated transformation (with reference to Potrykus, 1991). The employed transformation method is therefore restricted to microinjection of DNA into the isolated egg cell.

Agrobacterium-mediated transformation of isolated zygotes may appear to be hardly to manage, since there have so far been some unresolved technical requirements: The active release of signal molecules from the feeder-system which is indispensable for zygote survival and development has to be ensured throughout co-culture of zygotes and Agrobacteria and thereafter. While the zygotes are infected with Agrobacteria, the Agrobacteria should be kept apart from the co-cultured feeder cells. On the other hand, signal molecules from the feeder tissue have to have access to the isolated zygotes. Moreover, washing off the Agrobacteria from the extremely sensitive zygotes or zygote-derived structures and the transfer of the target cells to fresh medium after co-culture would undoubtedly cause severe cell damage or even collapse of the plasma membrane. After co-culture, these objects need to be transferred to an Agrobacterium-free medium which has exactly the same osmolality as the co-culture medium used before. Since the osmolality of the medium depends on the uptake of nutrients and the release of substances from the cells, after a few days practically every culture has its particular osmolality which is hardly reproducible in another medium containing actively developing feeder cells.

Despite the fact that wheat is the most widely-grown cereal crop in the world, unfortunately no efficient, cost-effective and genotype-independent method exists on the use of Agrobacterium transformation to generate stable, fertile transgenic wheat plants. A need therefore exists for an efficient method to genetically transform wheat genotypes of choice, preferentially elite breeding lines, the transgenic plants being uniformly transformed with a foreign gene. A need also exists for an efficient method for the production of transformed wheat plants capable of transmitting a foreign gene to progeny. In addition, a need exists for a method for Agrobacterium-mediated transformation of isolated wheat zygotes and subsequent generation of transgenic wheat plants. An additional need (especially in the light of regulatory requirements) exists for a transformation method which is independent of the use of a selectable marker gene (like e.g., antibiotic or herbicide resistance). Further, a technical solution is required to prevent the co-cultured feeder cells from being infected during co-culture of zygotes and Agrobacteria as well as to ensure transfer of the zygotes or zygote-derived structures to Agro-

bacterium-free, equiosmolar medium after co-culture.

Therefore, it is an objective of the current invention to provide a genotype-independent, efficient method for generating fertile, transgenic wheat plants.

## SUMMARY OF THE INVENTION

It is a particular object of the present invention to provide techniques that will allow to prepare transgenic, fertile Gramineae plants, which have been stably transformed through the introduction of one or more desired genes into the genome of these species. Accordingly a first embodiment of the invention is related to a method for producing a transgenic Gramineae plant comprising the steps of:

- (a) isolating a zygote from a Gramineae plant to be transformed in a way that said isolated zygote becomes substantially free from its naturally surrounding tissue,
- (b) introducing a DNA composition comprising a genetic component into the genome of said Gramineae plant, wherein said introduction is mediated by Agrobacterium transformation into said isolated zygote;
- (c) regenerating Gramineae plants from said zygotes which have received said genetic component; and
- (d) identifying a fertile, transgenic Gramineae plant whose genome has been altered through the stable introduction of said genetic component.

In a preferred embodiment, said Gramineae plant is selected from the group consisting of wheat, maize and barley. More preferably, the Gramineae plant is a Triticum species.

In another preferred embodiment, the regeneration step (c) of the method of the invention comprises a step, wherein the Gramineae plant is regenerated from said isolated zygote by a method comprising co-cultivation of said isolated zygote and/or zygotic embryo derived therefrom with dissected pistils, a cell culture derived from dissected pistils, or a culture comprising cells derived from dissected pistils. Preferably, the co-cultivation is carried out using a culture of immature wheat or barley pistils dissected at the one- or two-celled pollen stage of the floret.

In another preferred embodiment, the regeneration step (c) of the method of the invention comprises a step, wherein the Gramineae plant is regenerated from said isolated zygote by a method comprising co-cultivation of said isolated zygote and/or zygotic embryo derived therefrom with immature pollen, a cell culture derived from immature pollen, or a culture comprising cells derived from immature pollen. Preferably, the co-cultivation is carried out using a culture of isolated, androgenetically developing immature barley pollen.

In another preferred embodiment of the invention, the genetic component is transmitted through a complete sexual cycle of said transgenic Gramineae plant to its progeny.

Because of the high efficiency of the method of the invention, no selection or screening step based on selectable or screenable markers is necessary. The identification of transgenic Gramineae plants obtained by the method of the invention can be carried out simply by screening for the genetic component introduced into the genome of said Gramineae plant (e.g., by means of PCR or Southern blotting). Therefore, in another preferred embodiment, the transgenic Gramineae plant obtained by the method of the invention and/or the progeny thereof does not comprise a selectable or screenable marker gene.

In another preferred embodiment of the invention, the method does not comprise any step which leads to dedifferentiation of the zygote or the zygote-derived embryo. More preferably, no auxine or cytokinin compounds are utilized in a concentration and/or combination which leads to dedifferentiation.

In another preferred embodiment the genetic component introduced into the genome of said Gramineae plant comprises an expression cassette comprising a nucleic acid sequence operably linked to a promoter active in said Gramineae plant, wherein expression of said nucleic acid sequence confers a phenotypically distinguishable trait to said Gramineae plant.

## BRIEF DESCRIPTION OF THE DRAWINGS

The drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to these drawings in combination with the detailed description of specific embodiments presented herein.

Fig. 1: A: Millicell insert;  
B: semi-permeable membrane;  
C: isolated zygotes (or co-culture of zygotes and Agrobacteria)  
D: feeder system (e.g., androgenically developing pollen culture of barley)

Fig. 2: Plasmid map of pUGAB7 (binary vector comprising  $\beta$ -Glucuronidase encoding sequence and control of the maize ubiquitin promoter). PUBi-int: Maize ubiquitin 1 promoter comprising intron 1; GUS:  $\beta$ -Glucuronidase encoding sequence; P35S: CaMV 35S promoter; pat: phosphinotricinacetyltransferase (BASTA™ resistance gene); T35S: terminator sequence of CaMV 35S gene; Tnos: NOS terminator; LB/RB: left/right border of Agrobacterium T-DNA; Sm/Sp: Spectinomycin resistance gene; colE1: origin of replication; pVS1 ORI: origin of replication. Important restriction enzyme sites are indicated by name of the corresponding enzyme and the restriction site by base pair number.

Fig. 3: Plasmid map of pSUN-Ubi-GFP (binary vector comprising green fluorescence protein encoding sequence and control of the maize ubiquitin promoter). Ubi-1: Maize ubiquitin 1 promoter; GFP: green fluorescence protein encoding sequence; pat: phosphinotricinacetyltransferase (BASTA™ resistance gene); 35S-T: terminator sequence of CaMV 35S gene; nos-T: NOS terminator; LB/RB: left/right border of Agrobacterium T-DNA; aadA: Spectinomycin resistance gene; colE1: origin of replication. Important restriction enzyme sites are indi-

cated by name of the corresponding enzyme.

Fig. 4: Plasmid map of pSUN-Ubi-GUS (binary vector comprising  $\beta$ -Glucuronidase encoding sequence und control of the maize ubiquitin promoter). Ubi-1: Maize ubiquitin 1 promoter; GUS:  $\beta$ -Glucuronidase encoding sequence; pat: phosphotricinacetyltransferase (BASTA™ resistance gene); 35S-T: terminator sequence of CaMV 35S gene; nos-T: NOS terminator; LB/RB: left/right border of Agrobacterium T-DNA; aadA: Spectinomycin resistance gene; colE1: origin of replication. Important restriction enzyme sites are indicated by name of the corresponding enzyme.

Fig. 5: Transgene expression in an isolated wheat zygote after co-culture with LBA4404/ pSB1/ pYF133.

A: Two isolated zygotes shown in white light,

B: The same object under UV-light reveals *gfp* expression in one of the zygotes whereas the other zygote appears not to be transformed

#### GENERAL DEFINITIONS

It is to be understood that this invention is not limited to the particular methodology, protocols, cell lines, plant species or genera, constructs, and reagents described as such. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims. It must be noted that as used herein and in the appended claims, the singular forms "a," "and," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, reference to "a vector" is a reference to one or more vectors and includes equivalents thereof known to those skilled in the art, and so forth.

The term "about" is used herein to mean approximately, roughly, around, or in the region of. When the term "about" is used in conjunction with a numerical range, it modifies that range by extending the boundaries above and below the numerical values set forth. In general, the term "about" is used herein to modify a numerical value above and below the stated value by a variance of 20 percent up or down (higher or lower).

As used herein, the word "or" means any one member of a particular list and also includes any combination of members of that list.

The term "nucleic acid" refers to deoxyribonucleotides or ribonucleotides and polymers or hybrids thereof in either single-or double-stranded, sense or antisense form.

Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e. g., degenerate codon substitutions) and complementary sequences, as well as the sequence explicitly indicated. The term "nucleic acid" is used interchangeably herein with "gene", "cDNA", "mRNA", "oligonucleotide," and "polynucleotide".

The phrase "nucleic acid sequence" as used herein refers to a consecutive list of abbreviations, letters, characters or words, which represent nucleotides. In one embodi-



ment, a nucleic acid can be a "probe" which is a relatively short nucleic acid, usually less than 100 nucleotides in length. Often a nucleic acid probe is from about 50 nucleotides in length to about 10 nucleotides in length. A "target region" of a nucleic acid is a portion of a nucleic acid that is identified to be of interest. A "coding region" of a nucleic acid is the portion of the nucleic acid which is transcribed and translated in a sequence-specific manner to produce into a particular polypeptide or protein when placed under the control of appropriate regulatory sequences. The coding region is said to encode such a polypeptide or protein.

The term "antisense" is understood to mean a nucleic acid having a sequence complementary to a target sequence, for example a messenger RNA (mRNA) sequence the blocking of whose expression is sought to be initiated by hybridization with the target sequence.

The term "sense" is understood to mean a nucleic acid having a sequence which is homologous or identical to a target sequence, for example a sequence which binds to a protein transcription factor and which is involved in the expression of a given gene. According to a preferred embodiment, the nucleic acid comprises a gene of interest and elements allowing the expression of the said gene of interest.

The term "gene" refers to a coding region operably joined to appropriate regulatory sequences capable of regulating the expression of the polypeptide in some manner. A gene includes untranslated regulatory regions of DNA (e. g., promoters, enhancers, repressors, etc.) preceding (upstream) and following (downstream) the coding region (open reading frame, ORF) as well as, where applicable, intervening sequences (i.e., introns) between individual coding regions (i.e., exons).

As used herein the term "coding region" when used in reference to a structural gene refers to the nucleotide sequences which encode the amino acids found in the nascent polypeptide as a result of translation of a mRNA molecule. The coding region is bounded, in eukaryotes, on the 5'-side by the nucleotide triplet "ATG" which encodes the initiator methionine and on the 3'-side by one of the three triplets which specify stop codons (i.e., TAA, TAG, TGA). In addition to containing introns, genomic forms of a gene may also include sequences located on both the 5'- and 3'-end of the sequences which are present on the RNA transcript. These sequences are referred to as "flanking" sequences or regions (these flanking sequences are located 5' or 3' to the non-translated sequences present on the mRNA transcript). The 5'-flanking region may contain regulatory sequences such as promoters and enhancers which control or influence the transcription of the gene. The 3'-flanking region may contain sequences which direct the termination of transcription, posttranscriptional cleavage and polyadenylation.

The terms "polypeptide", "peptide", "oligopeptide", "polypeptide", "gene product", "expression product" and "protein" are used interchangeably herein to refer to a polymer or oligomer of consecutive amino acid residues.

The term "isolated" as used herein means that a material has been removed from its original environment. For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides can be part of a vector and/or such polynucleotides or polypep-



tides could be part of a composition, and would be isolated in that such a vector or composition is not part of its original environment.

5 The term "wild-type", "natural" or of "natural origin" means with respect to an organism, polypeptide, or nucleic acid sequence, that said organism is naturally occurring or available in at least one naturally occurring organism which is not changed, mutated, or otherwise manipulated by man.

10 The term "transgenic" or "recombinant" as used herein (e.g., with regard to a wheat zygote or wheat plants) is intended to refer to cells and/or plants that have incorporated exogenous genes or DNA sequences, including but not limited to genes or DNA sequences which are perhaps not normally present, genes not normally transcribed and translated ("expressed") in a given cell type, or any other genes or DNA sequences which one desires to introduce into the non-transformed cell and/or plant, such as  
15 genes which may normally be present in the non-transformed cell and/or plant but which one desires to have altered expression.

Preferably, the terms "transgenic" or "recombinant" with respect to, for example, a nucleic acid sequence (or an organism, expression cassette or vector comprising said  
20 nucleic acid sequence) refers to all those constructs originating by recombinant methods in which either

- a) said nucleic acid sequence, or
- b) a genetic control sequence linked operably to said nucleic acid sequence a), for example a promoter, or  
25 c) (a) and (b)

is not located in its natural genetic environment or has been modified by recombinant methods, an example of a modification being a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues. Natural genetic environment refers  
30 to the natural chromosomal locus in the organism of origin, or to the presence in a genomic library. In the case of a genomic library, the natural genetic environment of the nucleic acid sequence is preferably retained, at least in part. The environment flanks the nucleic acid sequence at least at one side and has a sequence of at least 50 bp, preferably at least 500 bp, especially preferably at least 1000 bp, very especially preferably at least 5000 bp, in length. A naturally occurring expression cassette - for example the naturally occurring combination of a promoter with the corresponding gene - becomes a recombinant expression cassette when it is modified by non-natural, synthetic "artificial" methods such as, for example, mutagenization. Such methods have been described (US 5,565,350; WO 00/15815). Preferably, the term "recombinant" with  
40 respect to nucleic acids as used herein means that the nucleic acid is covalently joined and adjacent to a nucleic acid to which it is not adjacent in its natural environment. "Recombinant" polypeptides or proteins refer to polypeptides or proteins produced by recombinant DNA techniques, i. e., produced from cells transformed by an exogenous recombinant DNA construct encoding the desired polypeptide or protein. Recombinant  
45 nucleic acids and polypeptide may also comprise molecules which as such does not exist in nature but are modified, changed, mutated or otherwise manipulated by man.

A "recombinant polypeptide" is a non-naturally occurring polypeptide that differs in se-

quence from a naturally occurring polypeptide by at least one amino acid residue. Preferred methods for producing said recombinant polypeptide and/or nucleic acid may comprise directed or non-directed mutagenesis, DNA shuffling or other methods of recursive recombination.

5

The terms "heterologous nucleic acid sequence" or "heterologous DNA" are used interchangeably to refer to a nucleotide sequence which is ligated to a nucleic acid sequence to which it is not ligated in nature, or to which it is ligated at a different location in nature. Heterologous DNA is not endogenous to the cell into which it is introduced, but has been obtained from another cell. Generally, although not necessarily, such heterologous DNA encodes RNA and proteins that are not normally produced by the cell into which it is expressed.

10

The "efficiency of transformation" or "frequency of transformation" as used herein can be measured by the number of transformed cells (or transgenic organisms grown from individual transformed cells) that are recovered under standard experimental conditions (i.e. standardized or normalized with respect to amount of cells contacted with foreign DNA, amount of delivered DNA, type and conditions of DNA delivery, general culture conditions etc.) For example, when isolated zygotes are used as starting material for transformation, the frequency of transformation can be expressed as the number of transgenic plant lines obtained per 100 isolated zygotes transformed.

15

20

The term "cell" refers to a single cell. The term "cells" refers to a population of cells. The population may be a pure population comprising one cell type. Likewise, the population may comprise more than one cell type. In the present invention, there is no limit on the number of cell types that a cell population may comprise. The cells may be synchronize or not synchronized, preferably the cells are synchronized.

25

The term "chromosomal DNA" or "chromosomal DNA-sequence" is to be understood as the genomic DNA of the cellular nucleus independent from the cell cycle status. Chromosomal DNA might therefore be organized in chromosomes or chromatids, they might be condensed or uncoiled. An insertion into the chromosomal DNA can be demonstrated and analyzed by various methods known in the art like e.g., PCR analysis, Southern blot analysis, fluorescence in situ hybridization (FISH), and in situ PCR.

30

35

An "embryo sac" is typically an eight-nucleate female gametophyte. The embryo sac arises from the megaspore by successive mitotic divisions.

40

A "megaspore" is one of the four haploid spores originating from the meiotic division of the diploid megaspore mother cell in the ovary and which gives rise to the megagametophyte.

A "microspore" is one of the four haploid spores originating from the meiotic division of the diploid microspore mother cell in the anther and which gives rise to the pollen grain.

45

The "polar nuclei" are the two nuclei of the central cell which occupies the central region of the embryo sac. Upon fusion of the central cell with one of the sperm cells, the polar nuclei unite with the sperm nucleus. The fertilized central cell constitutes the cellular origin of the endosperm.

5 The term "protoplasts" means protoplasts originating from any plant tissues where their entire wall is removed. They may be isolated from leaves, seeds, flowers, roots, pollen, embryos, ovaries or ovules or the protoplasts may be isolated from cell cultures including immature pollen, immature pollen derived embryos, zygotic embryos, secondary embryos from zygotic or immature pollen-derived embryos or immature pollen-derived cell-cultures. It is common knowledge how to isolate protoplasts by enzymatic removal of the cell wall. Techniques for this are described (in e.g. Potrykus 1986; Kranz 1991).

10 The term "structural gene" as used herein is intended to mean a DNA sequence that is transcribed into mRNA which is then translated into a sequence of amino acids characteristic of a specific polypeptide.

15 The term "expression" refers to the biosynthesis of a gene product. For example, in the case of a structural gene, expression involves transcription of the structural gene into mRNA and - optionally - the subsequent translation of mRNA into one or more polypeptides.

20 The term "transformation" includes introduction of genetic material into plant cells, preferably resulting in chromosomal integration and stable heritability through meiosis. Transformation also includes introduction of genetic material into plant cells in the form of plant viral vectors involving epichromosomal replication and gene expression which may exhibit variable properties with respect to meiotic stability.

25 The term "expression cassette" or "expression construct" as used herein is intended to mean the combination of any nucleic acid sequence to be expressed in operable linkage with a promoter sequence and - optionally - additional elements (like e.g., terminator and/or polyadenylation sequences) which facilitate expression of said nucleic acid sequence.

30 The term "promoter" as used herein is intended to mean a DNA sequence that directs the transcription of a DNA sequence (e.g., a structural gene). Typically, a promoter is located in the 5' region of a gene, proximal to the transcriptional start site of a structural gene. If a promoter is an inducible promoter, then the rate of transcription increases in response to an inducing agent. In contrast, the rate of transcription is not regulated by an inducing agent if the promoter is a constitutive promoter. Also, the promoter may be regulated in a tissue-specific or tissue preferred manner such that it is only active in transcribing the associated coding region in a specific tissue type(s) such as leaves, roots or meristem.

40 The term "operable linkage" or "operably linked" is to be understood as meaning, for example, the sequential arrangement of a regulatory element (e.g. a promoter) with a nucleic acid sequence to be expressed and, if appropriate, further regulatory elements (such as e.g., a terminator) in such a way that each of the regulatory elements can fulfill its intended function to allow, modify, facilitate or otherwise influence expression of said nucleic acid sequence. The expression may result depending on the arrangement of the nucleic acid sequences in relation to sense or antisense RNA. To this end, direct linkage in the chemical sense is not necessarily required. Genetic control sequences such as, for example, enhancer sequences, can also exert their function on the target

45

sequence from positions which are further away, or indeed from other DNA molecules. Preferred arrangements are those in which the nucleic acid sequence to be expressed recombinantly is positioned behind the sequence acting as promoter, so that the two sequences are linked covalently to each other. The distance between the promoter sequence and the nucleic acid sequence to be expressed recombinantly is preferably less than 200 base pairs, especially preferably less than 100 base pairs, very especially preferably less than 50 base pairs. Operable linkage, and an expression cassette, can be generated by means of customary recombination and cloning techniques as described (e.g., in Maniatis 1989; Silhavy 1984; Ausubel 1987; Gelvin 1990). However, further sequences which, for example, act as a linker with specific cleavage sites for restriction enzymes, or as a signal peptide, may also be positioned between the two sequences. The insertion of sequences may also lead to the expression of fusion proteins. Preferably, the expression cassette, consisting of a linkage of promoter and nucleic acid sequence to be expressed, can exist in a vector-integrated form and be inserted into a plant genome, for example by transformation.

The term "androgenetically developing pollen" is intended to mean any deviation from the designated development of a microspore into a functional pollen grain. In contrast to the normal pollen formation, androgenetically developing pollen undergoes a number of successive cell divisions to form non-embryogenic or embryogenic callus or an embryo-like structure. These structures which genetically correspond to the male gametophyte can develop into haploid plants. The development of these structures may also include spontaneous or artificially induced genome doubling which may result in regeneration of homozygous, fertile plants.

## DETAILED DESCRIPTION OF THE INVENTION

The method according to the invention is remarkable in that the transformed zygotes are able without additional treatment by e.g., hormones to regenerate into green fertile plants. Furthermore it is not necessary to introduce a selectable gene. The present invention has the advantage of being genotype-independent. Since zygote development is a universal step in higher plants and in particular within the Gramineae family, the methods disclosed herein can be transferred at least to other members of said family.

The terms "Gramineae" or "Graminaceae" as used herein intends to comprise all plants species of the Gramineae (Poaceae) family, especially those employed as foodstuffs or feeding stuffs such as rice, maize, wheat or other cereal species such as barley, millet and sorghum, rye, triticale or oats, and sugar cane, and all grass species. Furthermore included are the mature plants, seed, shoots and seedlings, and parts, propagation material and cultures derived therefrom, for example cell cultures. Mature plants refers to plants at any developmental stage beyond that of the seedling. The term seedling refers to a young immature plant in an early developmental stage, at which it is still dependent upon assimilates stored within the seed (e.g. in the endosperm, perisperm or cotyledons).

Included are all genera of the subfamilies Bambusoideae (e.g., the genus bamboo), Andropogonoideae (e.g., the genera Saccharum, Sorghum, or Zea), Arundineae (e.g.,

the genus *Phragmites*), *Oryzoideae* (e.g., the genus *Oryza*), *Panicoideae* (e.g., the genera *Panicum*, *Pennisetum*, and *Setaria*), *Pooideae* (*Festuciadeae*) (e.g., the genera *Poa*, *Festuca*, *Lolium*, *Trisetum*, *Agrostis*, *Phleum*, *Dactylis*, *Alopecurus*, *Avena*, *Triticum*, *Secale*, and *Hordeum*).

Preferred are *Avena sativa* (oats), *Bambusa* sp. and *Bambusa bambos* (bamboo), *Saccharum officinarum* (sugar cane), *Triticum dicoccum* (Emmer wheat), *Triticum monococcum* (Einkorn wheat), *Triticum spelta* (spelt wheat), *Triticum durum* (wheat), *Triticum turgidum*, *Triticum aestivum* (wheat), *Zea mays* (maize/corn), *Panicum miliaceum* (common millet), *Pennisetum thipoides* (Bulrush millet), *Hordeum vulgare* or *H. sativum* (barley), *Oryza sativa* (rice), *Zizania aquatica* (wild rice), *Secale cereale* (rye), *Sorghum bicolor* (*S. vulgare*) (sorghum). More preferred are wheat species and species with nearly identical zygote development characteristics like barley or maize. Especially preferred are all wheat species especially of the *Triticum* family (including both winter and spring wheat), more especially *Triticum* spp.: common (*T. aestivum*), durum (*T. durum*), spelt (*T. spelta*), *Triticum dicoccum* (Emmer wheat), *Triticum turgidum*, and *Triticum monococcum* (Einkorn wheat), with *T. aestivum* being particularly preferred. The method of the invention can be used to produce transgenic plants from spring wheats, such as, for example, Bobwhite, Marshall, PIVOT1, UC702, and Panewawa as well as from winter wheats, such as, for example, HY368, Neeley, FL302, RH91, R332, R1269 and R585. However, it should be pointed out, that the method of the invention is not limited to certain varieties but is highly genotype-independent.

## 25 ZYGOTE ISOLATION

Isolated zygotes as used within the method of the invention have demonstrated a high potential for regeneration.

The term "isolated zygote" as used herein means the fertilized egg cell of a Gramineae plant substantially free of the surrounding tissue (like, e.g. pericarp, the two integuments, nucellus cells, embryo sac, constituting cells of the embryo sac).

The zygote in higher plants is per definition totipotent, i.e., its natural fate (function) is to develop into an intact plant.

The ovule is located inside the ovary and consists of the two integuments, nucellus tissue and the embryo sac. The embryo sac consists of a large central cell with two central or polar nuclei. At the chalazal end a number of antipodal cells are present while the two synergids and the egg cell are located within the micropylar region of the embryo sac. The structure of the embryo sac before, during and after fertilization has been described (Mogensen 1982; Engel 1989). In barley and wheat, fertilization occurs within about one hour after pollination: The pollen tube grows through the stigmata towards the ovule, then along the integuments, and eventually enters the micropyle and one of the synergids, which at this time has degenerated. There, the two sperm cells are released from the pollen tube, and one fuses with the egg while the other one fuses with the central cell. The fertilized, triploid central cell will at a later stage give rise to the triploid endosperm. Within about one day, wheat and barley zygotes have completed their first division which results in the two-celled proembryo.

Different methods for the isolation of Gramineae zygotes have been described (Holm et al. 1994, Leduc et. al. 1996, Kumlehn et al. 1998; see also Example 1). The unfertilized egg cell is a true protoplast. However, since immediately after fertilization cell wall formation starts, the zygote is by definition not a protoplast. However, may features of a protoplast still exist.

The behavior of the zygote is to some extent depending on the osmotic pressure of the medium. The osmotic pressure of the medium thus has to be of an order of magnitude where the zygote neither is swelling to an extent that it ruptures or undergoes extensive shriveling due to dehydration. Osmotic pressures ranging from 250 to 750 mOsm/kg can be tolerated, in particular a pressure of 375 mOsm/kg is suitable for barley zygote isolation, and an osmolality of about 600 mOsm/kg for wheat zygote isolation. The osmotic pressure can be increased or reduced by addition/removal of e.g. mannitol. For example, a 0.55 M mannitol solution guarantees an appropriate osmotic pressure for wheat zygote isolation. During isolation, the zygotes may become sticky in the presence of exogenously supplied  $\text{Ca}^{2+}$  making the cells very difficult to handle. Therefore, in a preferred embodiment pure mannitol solution is used in which the isolated zygotes were morphologically stable and did not show any obvious loss of viability during preparation.

With regard to wheat, efficient and reproducible embryo development has been obtained from fertilized wheat (*Triticum aestivum* L.) egg cells isolated 3-6 h after hand-pollination of emasculated spikes. It is possible to routinely isolate viable zygotes from about 75% of the processed pistils from cultivars of both winter and spring type.

#### AGROBACTERIUM TRANSFORMATION

The genetic component is introduced into the isolated zygote by means of Agrobacterium-mediated DNA transfer.

The term "Agrobacterium" as used herein means all species of the Agrobacterium family (including Agrobacterium tumefaciens and Agrobacterium rhizogenes). Preferably, transformation is realized utilizing strains of Agrobacterium tumefaciens or Agrobacterium rhizogenes. The principles of plant transformation by means of Agrobacterium-mediated DNA transfer are well known in the art (see above; BACKGROUND OF THE INVENTION; Horsch 1985).

Agrobacterium tumefaciens and A. rhizogenes are plant-pathogenic soil bacteria, which genetically transform plant cells. The Ti and Ri plasmids of A. tumefaciens and A. rhizogenes, respectively, carry genes responsible for genetic transformation of the plant (Kado 1991). Vectors are based on the Agrobacterium Ti- or Ri-plasmid and utilize a natural system of DNA transfer into the plant genome. As part of this highly developed parasitism Agrobacterium transfers a defined part of its genomic information (the T-DNA; flanked by about 25 bp repeats, named left and right border) into the chromosomal DNA of the plant cell (Zupan 2000). By combined action of the so called vir genes (part of the original Ti-plasmids) said DNA-transfer is mediated. For utilization of this natural system, Ti-plasmids were developed which lack the original tumor inducing genes ("disarmed vectors"). In a further improvement, the so called "binary vector

systems", the T-DNA was physically separated from the other functional elements of the Ti-plasmid (e.g., the *vir* genes), by being incorporated into a shuttle vector, which allowed easier handling (EP-A 120 516; US 4.940.838). These binary vectors comprise (beside the disarmed T-DNA with its border sequences), prokaryotic sequences for replication both in *Agrobacterium* and *E. coli*. It is an advantage of *Agrobacterium*-mediated transformation that in general only the DNA flanked by the borders is transferred into the genome and that preferentially only one copy is inserted. Descriptions of *Agrobacterium* vector systems and methods for *Agrobacterium*-mediated gene transfer are known in the art (Miki 1993; Gruber 1993; Moloney 1989).

Hence, for *Agrobacteria*-mediated transformation the genetic composition (e.g., comprising an expression cassette) is integrated into specific plasmids, either into a shuttle or intermediate vector, or into a binary vector. If a Ti or Ri plasmid is to be used for the transformation, at least the right border, but in most cases the right and left border, of the Ti or Ri plasmid T-DNA is linked to the expression cassette to be introduced in the form of a flanking region. Binary vectors are preferably used. Binary vectors are capable of replication both in *E.coli* and in *Agrobacterium*. They may comprise a selection marker gene and a linker or polylinker (for insertion of e.g. the expression cassette to be transferred) flanked by the right and left T-DNA border sequence. They can be transferred directly into *Agrobacterium* (Holsters 1978). The selection marker gene permits the selection of transformed *Agrobacteria* and is, for example, the *nptII* gene, which confers resistance to kanamycin. The *Agrobacterium* which acts as host organism in this case should already contain a plasmid with the *vir* region. The latter is required for transferring the T-DNA to the plant cell. An *Agrobacterium* transformed in this way can be used for transforming plant cells. The use of T-DNA for transforming plant cells has been studied and described intensively (EP 120 516; Hoekema 1985; An et al. 1985).

Common binary vectors are based on "broad host range"-plasmids like pRK252 (Bevan 1984) or pTJS75 (Watson et al. 1985) derived from the P-type plasmid RK2. Most of these vectors are derivatives of pBIN19 (Bevan 1984). Various binary vectors are known, some of which are commercially available such as, for example, pBI101.2 or pBIN19 (Clontech Laboratories, Inc. USA). Additional vectors were improved with regard to size and handling (e.g. pPZP; Hajdukiewicz 1994). Improved vector systems are described also in WO 02/00900.

In a preferred embodiment, *Agrobacterium* strains for use in the practice of the invention include octopine strains, e.g., LBA4404 or agropine strains, e.g., EHA101 or EHA105. Suitable strains of *A. tumefaciens* for DNA transfer are for example EHA101pEHA101 (Hood et al. 1986), EHA105[pEHA105] (Li 1992), LBA4404[pAL4404] (Hoekema 1983), C58C1[pMP90] (Koncz & Schell 1986), and C58C1[pGV2260] (Deblaere 1985). Other suitable strains are *Agrobacterium tumefaciens* C58, a nopaline strain. Other suitable strains are *A. tumefaciens* C58C1 (Van Larebeke 1974), A136 (Watson 1975) or LBA4011 (Klapwijk 1980). In a preferred embodiment, the *Agrobacterium* strain used to transform the plant tissue pre-cultured with the plant phenolic compound contains a L,L-succinamopine type Ti-plasmid, preferably disarmed, such as pEHA101. In another preferred embodiment, the *Agrobacterium* strain used to transform the plant tissue pre-cultured with the plant phenolic compound contains an octopine-type Ti-plasmid, preferably disarmed, such as pAL4404. Gener-



ally, when using octopine-type Ti-plasmids or helper plasmids, it is preferred that the virF gene be deleted or inactivated (Jarschow 1991).

5 The method of the invention can also be used in combination with particular Agrobacterium strains, to further increase the transformation efficiency, such as Agrobacterium strains wherein the vir gene expression and/or induction thereof is altered due to the presence of mutant or chimeric virA or virG genes (e.g. Hansen 1994; Chen and Wijnans 1991; Scheeren-Groot et al., 1994).

10 Preferred are further combinations of Agrobacterium tumefaciens strain LBA4404 (Hiei 1994) with super-virulent plasmids. These are preferably pTOK246-based vectors (Ishida 1996).

15 A binary vector or any other vector can be modified by common DNA recombination techniques, multiplied in *E. coli*, and introduced into Agrobacterium by e.g., electroporation or other transformation techniques (Mozo 1991).

20 Agrobacterium is grown and used in a manner similar to that described in Ishida (Ishida et al. 1996). The vector comprising Agrobacterium strain may, for example, be grown for 3 days on YP medium (5 g/l yeast extract, 10 g/l peptone, 5 g/l Nail, 15 g/l agar, pH 6.8) supplemented with the appropriate antibiotic (e.g., 50 mg/l spectinomycin). Bacteria are collected with a loop from the solid medium and resuspended. In a preferred embodiment of the invention, Agrobacterium cultures are started by use of aliquots frozen at -80°C. For Agrobacterium treatment of isolated zygotes, the bacteria are  
25 resuspended in the medium used for zygote culture.

The zygote to be transformed may be treated with cell wall degrading enzymes to facilitate or enhance transformation efficacy. However, especially for wheat zygotes isolated within 20 hours after pollination this degradation step is not necessary and is omitted in  
30 a preferred embodiment of the invention. The concentration of Agrobacterium used for infection and co-cultivation may need to be varied. It is understood in the art that high densities and long co-cultivation can injure the isolated zygotes. Thus, a range of Agrobacterium concentrations from  $10^5$  -  $10^{10}$  cfu/ml and a range of co-cultivation periods from a few hours to 7 days are tested with the isolated zygotes. The co-cultivation  
35 of Agrobacterium with the isolated zygotes is in general carried out for 1 to five, preferably 2 to 3 days. In an improved embodiment of the invention the isolated zygotes may be treated with a phenolic compound prior to or during the Agrobacterium co-cultivation. "Plant phenolic compounds" or "plant phenolics" suitable within the scope of the invention are those isolated substituted phenolic molecules which are capable to  
40 induce a positive chemotactic response, particularly those who are capable to induce increased vir gene expression in a Ti-plasmid containing Agrobacterium sp., particularly a Ti-plasmid containing Agrobacterium tumefaciens. Methods to measure chemotactic responses towards plant phenolic compounds have been like e.g., described (Ashby 1988) and methods to measure induction of vir gene expression are also well  
45 known (Stachel 1985; Bolton 1986). The pre-treatment and/or treatment during Agrobacterium co-cultivation has at least two beneficial effects: Induction of the vir genes of Ti plasmids or helper plasmids (Van Wordragen 1992; Jacq 1993; James 1993; Guivarc'h 1993), and enhancement of the competence for incorporation of foreign DNA into the genome of the plant cell.

Preferred plant phenolic compounds are those found in wound exudates of plant cells. One of the best known plant phenolic compounds is acetosyringone, which is present in a number of wounded and intact cells of various plants, albeit in different concentrations. However, acetosyringone (3,5-dimethoxy-4-hydroxyacetophenone) is not the only plant phenolic which can induce the expression of vir genes. Other examples are  $\alpha$ -hydroxy-acetosyringone, sinapinic acid (3,5-dimethoxy-4-hydroxycinnamic acid), syringic acid (4-hydroxy-3,5 dimethoxybenzoic acid), ferulic acid (4-hydroxy-3-methoxycinnamic acid), catechol (1,2-dihydroxybenzene), p-hydroxybenzoic acid (4-hydroxybenzoic acid),  $\beta$ -resorcylic acid (2,4-dihydroxybenzoic acid), protocatechuic acid (3,4-dihydroxybenzoic acid), pyrogalllic acid (2,3,4-trihydroxybenzoic acid), gallic acid (3,4,5-trihydroxybenzoic acid) and vanillin (3-methoxy-4-hydroxybenzaldehyde), and these phenolic compounds are known or expected to be able to replace acetosyringone in the cultivation media with similar results. As used herein, the mentioned molecules are referred to as plant phenolic compounds.

Plant phenolic compounds can be added to the plant culture medium either alone or in combination with other plant phenolic compounds. A particularly preferred combination of plant phenolic compounds comprises at least acetosyringone and p-hydroxybenzoic acid; but it is expected that other combinations of two, or more, plant phenolic compounds will also act synergistically in enhancing the transformation efficiency.

Moreover, certain compounds, such as osmoprotectants (e.g. L-proline preferably at a concentration of about 700 mg/L or betaine), phytohormes (inter alia NAA), opines, or sugars, are expected to act synergistically when added in combination with plant phenolic compounds.

In one embodiment of the invention, it is preferred that the plant phenolic compound, particularly acetosyringone is added to the medium prior to contacting the isolated zygotes with *Agrobacteria* (for e.g., several hours to one day). The exact period in which the cultured cells are incubated in the medium containing the plant phenolic compound such as acetosyringone, is believed not to be critical and only limited by the time the zygote starts entering mitosis.

The concentration of the plant phenolic compound in the medium is also believed to have an effect on the development of competence for integrative transformation. The optimal concentration range of plant phenolic compounds in the medium may vary depending on the species from which the zygote derived, but it is expected that about 100  $\mu$ M to 700  $\mu$ M is a suitable concentration for many purposes. However, concentrations as low as approximately 25  $\mu$ M can be used to obtain a good effect on transformation efficiency. Likewise, it is expected that higher concentrations up to approximately 1000  $\mu$ M will yield similar effects. Comparable concentrations apply to other plant phenolic compounds, and optimal concentrations can be established easily by experimentation in accordance with this invention.

*Agrobacteria* to be co-cultivated with the isolated zygotes can be either pre-incubated with acetosyringone or another plant phenolic compound, as known by the person skilled in the art, or used directly after isolation from their culture medium. Particularly suited induction conditions for *Agrobacterium tumefaciens* have been described by

Vernade et al. (1988). Efficiency of transformation with *Agrobacterium* can be enhanced by numerous other methods known in the art like for example vacuum infiltration (WO 00/58484), heat shock and/or centrifugation, addition of silver nitrate, sonication etc.

5       Supplementation of the co-culture medium with antioxidants (e.g., dithiothreitol), phenol-absorbing compounds (like polyvinylpyrrolidone, Perl et al. 1996) or thiol compounds (e.g., L-cysteine, Olhoft et al. 2001) which can decrease tissue necrosis due to plant defence responses (like phenolic oxidation) may further improve the efficiency of  
10       *Agrobacterium*-mediated transformation.

It has been observed within this invention that transformation efficacy of the isolated zygotes by *Agrobacterium* can be significantly improved by keeping the pH of the co-cultivation medium in a range from 5.4 to 6.4, preferably 5.6 to 6.2, especially preferably  
15       5.8 to 6.0. In an improved embodiment of the invention stabilisation of the pH in this range is mediated by a combination of MES and potassium hydrogenphosphate buffers.

#### CULTIVATION OF ZYGOTES AND REGENERATION INTO PLANTS

20       After transformation, the zygote is transferred into another Millicell insert placed in the same petri dish and the co-culture insert is removed to avoid an ongoing impact of the *Agrobacteria* on the medium (nutrient depletion, release of bacterial exudates, modification of pH). It turned out to be crucial to keep the isolated zygotes or zygote-derived  
25       structures within the same medium, because even a minor shift in concentration of some media components or in osmolality causes subsequent degeneration of the cultured cells. Since the medium used for co-culture is basically suitable for embryo development, only some supplementation appeared to be necessary, e.g. antibiotics (like  
30       e.g., Amoxicillin, Cefotaxime, Clavulanic acid, Timentin or a combination thereof) and additional  $\text{CaCl}_2$  to stop the development of the *Agrobacteria* as well as additional glutamine to support embryonic development.

In a preferred embodiment of the invention, the in vitro development of the zygotes and their subsequent regeneration into fertile plants is mediated by co-cultivation techniques. The co-cultivation principle is well established in tissue culture, where e.g., protoplasts often are co-cultivated with suspension cultures to support protoplast development (Evans 1983). There is today very little evidence on the nature of the factors which during the co-cultivation support regeneration and embryogenesis. Methods for  
35       co-cultivation to be preferably utilized herein are described (e.g., in WO 94/01999; Kümlehn et al. 1998).  
40

Several kinds of cells and/or tissues are suitable for regeneration of the zygotes into embryos. Within this invention, these cells and/or tissues are subsumed under the term "feeder system". As feeder systems, embryogenic as well as non-embryogenic  
45       suspensions can be used. Preferably, a co-cultivation with isolated immature pollen (preferably barley pollen) or isolated immature pollen which had been stimulated to androgenetic development is used. Alternatively, cultures of wheat pistils (gynoecia) can be employed for feeder systems. Methods to co-culture ovaries are known in the art (Köhler and Wenzel 1978, Mejza et al. 1993). Furthermore, a combination of the above

mentioned feeder systems can be used like e.g., a combination of barley immature pollen culture with wheat pistil culture.

5 Thus, in another preferred embodiment, the regeneration step (c) of the method of the invention comprises a step, wherein the Gramineae plant is regenerated from said isolated zygote by a method comprising co-cultivation of said isolated zygote and/or zy-  
10 gotic embryo derived therefrom with dissected pistils, a cell culture derived from dissected pistils, or a culture comprising cells derived from dissected pistils. Preferably, the co-cultivation is carried out using a culture of immature wheat or barley pistils dissected at the one- or two-celled pollen stage of the floret.

Co-culture with e.g. barley pollen cultures resulted in embryonic development of the cultivated wheat zygotes. Within 23 h of pollination the zygotes underwent the first cell  
15 division and then formed differentiated embryos very similar to those grown in planta.

Pollen embryogenesis is one of the few tissue culture systems available where a very large number of cells simultaneously undergo embryonic development. Immature barley pollen with their high regeneration potential seem to condition the medium with a complex combination of compounds which support embryonic development of zygotes  
20 within the method of the invention.

This complex mixture of components may be continuously released when immature pollen undergoes embryogenesis. As an alternative to the co-cultivation procedure, it is thus possible, to use conditioned medium or extracts of immature pollen or anthers, in  
25 particular pollen, which is undergoing embryogenesis. Extracts are defined as non-intact cellular materials, which are isolated from pollen or anthers, f.ex. by blending and possibly further purified by filtration or fractionation.

In general, immature pollen or anthers from any species or variety may be used but the  
30 regeneration frequency of protoplasts, cells and tissues may depend on the source of the feeder tissue. The regeneration frequency also depends on the cell density of the pollen culture or the anthers cultivated in the medium. Immature pollen or anthers from barley (e.g., the barley cultivar Igri) are particularly suitable for use in the method according to the invention. In a preferred embodiment, immature pollen of cv. Igri are iso-  
35 lated and cultured as basically described by Mordhorst and Lörz (1993) and transferred to the medium used for zygote culture as an intermediate step.

It is preferred to use immature barley pollen cultures at a density between  $10^2$  to  $10^6$ , preferably  $10^3$  to  $3 \cdot 10^5$ , especially preferably  $10^4$  and  $10^5$  immature pollen per ml of  
40 culture medium or the equivalent amount of anthers. It is preferred, however, to use immature pollen. Suspension cultures or callus cultures derived from immature pollen can also be used (Roberts-Oehlschlager et al. 1990). It is preferred though to use young cultures of immature pollen or anthers, i.e., preferably cultures that are younger than 30 days, and in particular younger than 21 days, more preferably cultures be-  
45 tween 5 and 14 days.

Feeder system-mediated co-cultivation may involve a physical separation between the supporting feeder system and the isolated zygotes to be supported and where it is ensured that low and high molecular compounds can diffuse between the two cultures

5 The androgenetically developing pollen and the plant material to be regenerated can be cultured in the same medium which preferably is liquid or semi-liquid or the pollen-derived microcalli can be cultivated in one medium and the plant material to be regenerated in a second medium which is in exchangeable contact with the first medium or contain fractions of the first medium. This can, for example be realized by cultivating the isolated zygotes in cell culture dishes with bottom semi-permeable membranes (e.g. Millicell inserts; Millipore, Bedford, Mass., USA) placed in larger cell culture dishes comprising the pollen culture.

10 A co-cultivation system may of course be constructed in a number of ways. The immature pollen may be cultured directly in the petri dish thereby avoiding the use of a large insert. The pollen-derived microcalli and the isolated zygotes to be regenerated may also be present in the same reservoir or dish, thereby avoiding the use of any of the two inserts. The isolated zygotes to be regenerated may be embedded in a solid medium such as agarose which is permeable for the components present in the pollen culture. The only essential principle in the co-cultivation is thus that the components present in the pollen culture undergoing embryogenesis readily can reach the material (e.g., the zygotes) to be regenerated.

20 It was found that the number of zygotes, cultivated per Millicell insert did not influence the embryonic development that is, even a singly cultivated zygote can develop into a well-differentiated embryo (Kumlehn et al. 1998).

25 It is of course obvious that the growth and culture media used for the tissue culture contain the nutrients essential for the plant material. The composition of nutrients in such growth and culture media are common knowledge to any person experienced in the field (e.g., Kao 1975; Linsmaier 1965; Spencer 1990; Twell 1989a). The N6Z medium has demonstrated to be especially preferably for promoting embryogenesis from isolated wheat zygotes (Table 1).

30 The feeder system can be employed only after the Agrobacterium co-cultivation step but may be also – preferably – employed immediately after isolation of the zygote and remained during the Agrobacterium co-cultivation. In another preferred embodiment of the invention, the Agrobacteria are kept separate from the feeder system and only contact with the zygotes to be transformed is allowed. This may be ensured by incubating the zygotes with the Agrobacteria within a Millicell insert having a pore width of 0.4  $\mu\text{m}$  which prevents diffusion of Agrobacteria into the medium outside this insert as well as contamination of the feeder system. The feeder system may be cultured directly in a larger cell culture plate comprising the Millicell used for co-culture of the zygotes with Agrobacteria (Fig. 1). This approach is advantageous since contamination of the feeder system with Agrobacterium is prevented. This allows to use the very same feeder system also after removal of the Agrobacterium for further incubation/regeneration of the zygotes. Zygotes would be simply transferred to another Millicell insert and again cultivated within the before used feeder system comprising plate (supplemented with antibiotics and/or Calcium chloride to prevent growth of remaining Agrobacteria in the zygote culture). This variant is especially preferred in another improved embodiment of the invention, since it helps to overcome certain difficulties related to sensitivity of zygotes to changed environmental conditions (like e.g., changed medium composition,

osmolality and pH).

5 In a preferred embodiment, a co-culture with barley immature pollen which had been stimulated to androgenetic development is employed. This resulted in embryonic development of the cultivated wheat zygotes with high efficiency. Within 23 h of pollination, the zygotes underwent their first cell division. They proceeded to develop into club-shaped embryos, most of which turned subsequently to dorsoventral differentiation. The morphological patterns of in-vitro-grown embryos were in accordance with those of normal zygotic embryos growing in planta. Optimized media are disclosed in examples 4 and 5 (Table 1). More than 80% of the zygotes eventually developed into plants (e.g., in genotypes Florida and Veery #5, respectively). All regenerated plants were morphologically normal and fertile.

15 After development of zygote-derived embryos (preferably of a size of at least 1 mm; which is obtained for example after three to four weeks of co-cultivation with immature pollen in the dark at 25°C), these embryos may further be cultivated towards development of plantlets and plants. For this purpose numerous methods known to the person skilled in the art may be employed. For example, the zygote-derived embryos may be transferred (plated) e.g., on solidified medium (e.g., K4NT, Table 1) supporting further embryo differentiation, germination and plantlet formation. The medium used in this step can be any medium which permits germination of embryos. In one embodiment, a shoot-producing compound is added to the medium. These shoot-producing compounds are well known in the art (Mursahige and Skoog, 1962; Kasha 1990). Such compounds include plant growth regulators and include IAA, IBA, and BA at low concentrations (Becker *et al.*, 1994; Vasil 1992). In another embodiment of the invention, a medium free of a plant growth regulator can be used to induce shoot formation (Weeks 1993). Subculture can be carried out on a monthly basis until plantlets are transferred to soil.

30 If necessary, explants with shoots can be transferred to medium particularly supporting root formation. These media are well known in the art (Weeks 1993; Vasil 1992). One preferred root-producing medium is a modified MS-medium without any plant growth regulator (Murashige 1962; Zhou 1992). Once roots have been formed, the plants can be transferred to soil and grown following methods known in the art to produce mature plants with seeds.

40 In another embodiment of the invention, the media as employed during the method of the invention may be optionally further supplemented with one or more plant growth regulator, like e.g., cytokinin compounds (e.g., 6-Benzylaminopurin) or auxin compounds (e.g., 2,4-D). The term "plant growth regulator" (PGR) as used herein means naturally occurring or synthetic (not-naturally occurring) compounds that can regulate plant growth and development. PGRs may act singly or in consort with one another or with other compounds (e.g., sugars, amino acids).

45 The term "auxin" or "auxin compounds" comprises compounds which stimulate cellular elongation and division, differentiation of vascular tissue, fruit development, formation of adventitious roots, production of ethylene, and - in high concentrations - induce dedifferentiation (callus formation). The most common naturally occurring auxin is indoleacetic acid (IAA), which is transported polarly in roots and stems. Synthetic auxins

are used extensively in modern agriculture. Auxin compounds comprise indole-3-butyric acid (IBA), naphthylacetic acid (NAA), and 2,4-dichlorophenoxyacetic acid (2,4-D), wherein 2,4-D is a more active compound.

5 The term "cytokinin" or "cytokinin compound" comprises compounds which stimulate cellular division, expansion of cotyledons, and growth of lateral buds. They delay senescence of detached leaves and, in combination with auxines (e.g. IAA), may influence formation of roots and shoots. Cytokinin compounds comprise, for example, 6-isopentenyladenine (IPA) and 6-benzyladenine/6-benzylaminopurine (BAP).

10

In a preferred embodiment the method of the invention is carried out under conditions which does not lead to dedifferentiation. Preferably, no dedifferentiation is induced to the isolated zygote and zygote-derived embryo, plantlet or plant. More preferably, no embryonic tissue dedifferentiation and/or formation of callus or callus-like structures is initiated.

15

The term "dedifferentiation" as used herein is intended to mean the process of formation of rapidly dividing cells without particular function in the scope of the plant body. These cells often possess an increased potency with regard to its ability to develop into various plant tissues. Preferably the term is intended to mean the reversion of a differentiated or specialized tissues to a more pluripotent or totipotent (e.g., embryonic) form. Dedifferentiation may lead to reprogramming of a plant tissue (revert first to undifferentiated, non-specialized cells. then to new and different paths). The term "totipotency" as used herein is intended to mean a plant cell containing all the genetic and/or cellular information required to form an entire plant. Dedifferentiation can be initiated by certain plant growth regulators (e.g., auxin and/or cytokinin compounds), especially by certain combinations and/or concentrations thereof.

20

25

In a preferred embodiment, the concentration and/or combination of auxins and/or cytokinin added to a culture medium used within the method of the invention is kept in a range, where no dedifferentiation (e.g., callus induction) is initiated. Combinations of auxins/cytokinins leading to shoot and/or root induction may be advantageously employed during plant regeneration from the zygote-derived embryo.

30

35 Within an preferred embodiment of the method of the invention cytokinins and auxins may be utilized to induce shoot and/or root development, but not dedifferentiation. However, under certain circumstances, dedifferentiation of a transformed zygotic embryo in a later step of the method may be advantageous (e.g., to increase the probability of systemically (non-chimeric) transformation of the Gramineae plant obtained by the method of the invention or to multiply certain transformation events). In general, however, preferably no dedifferentiation is applied.

40

More preferably no auxin compound (like e.g., 2,4-D) are added or their concentration is kept in a low range of 0 to about 0.2 mg/l 2,4-D.

45

The media as used herein (e.g., for cultivation of zygotes or zygote-derived embryos) may be liquid media (e.g., for pollen and zygote cultures) or solid media (e.g., solidified by 2-7 g/l Gelrite® (Sigma Chemical, St. Louis, MO) or 4 g/l low-melting agarose).



Other important aspects of the invention include the progeny of the transgenic plants prepared by the disclosed methods, as well as the cells derived from such progeny, and the seeds obtained from such progeny.

- 5 Within this invention a highly efficient transformation and regeneration protocol for isolated Gramineae (especially wheat) zygotes undergoing direct embryo differentiation when co-cultured with immature pollen. Almost every wheat zygote can undergo embryonic development with morphological patterns homologous to normal zygotic embryos growing in planta. In this way, it is possible to regenerate plant materials, which
- 10 previously have been difficult to regenerate from cell cultures into green fertile plants in much higher frequencies than seen before and it is also possible to regenerate plant materials, which previously have been impossible to regenerate.

## SELECTION

5 The method of the invention reliably leads to transgenic, non-chimeric gramineae plants with high efficiency. Therefore, utilization of selectable or screenable marker genes for identification of the transgenic plants is not necessary. Because of the reservations of the public with regard to antibiotic and other selectable markers this is considered to be an economically highly relevant advantage over the methods known in the art.

10 The insertion of the genetic component into the chromosomal DNA can be demonstrated and analyzed by various methods known in the art like e.g., PCR analysis, Southern blot analysis, fluorescence in situ hybridization (FISH), and in situ PCR.

15 However, insertion of a selectable and/or screenable marker is comprised within the scope of the method of the invention. This may be advantageous e.g., for later use as a herbicide-resistance trait. Therefore, in another embodiment of the invention the genetic component introduced into the genome of the Gramineae plant comprises an expression cassette able to express a selectable or screenable marker in said Gramineae plant. The medium employed for culturing the transformed zygotes and/or zygote-  
20 derived embryos and/or plantlets may further contain a compound used to select zygotes, embryos or plantlets containing a selectable or screenable DNA sequence.

The compound used for said selection or screening can be any of a variety of well known selection compounds, such as antibiotics and herbicides. Preferred compounds  
25 can include geneticin (G-418) (aminoglycoside) (Nehra et al. 1994), glyphosate (Della-Cioppa et al. 1987) and bialaphos (Vasil et al. 1992; Weeks et al. 1993). The availability of alternative selection agents is an important requirement for commercial application of agriculture biotechnology. The use of kanamycin has been less successful for cereal crops because of the high endogenous level of tolerance (Dekeyser et al. 1989).  
30 Bialaphos has been widely used as a selection agent in cereal crop transformation (Weeks et al. 1993; Vasil et al. 1993; Becker et al. 1994; Nehra et al. 1994; Wan and Lemaux 1994). Further preferred selectable and screenable marker genes are disclosed below.

35 Transformed plant material (e.g., cells, embryos, tissues or plantlets) which express such marker genes are capable of developing in the presence of concentrations of a corresponding selection compound (e.g., antibiotic or herbicide) which suppresses growth of an untransformed wild type tissue. The resulting plants can be bred and hybridized in the customary fashion. Two or more generations should be grown in order  
40 to ensure that the genomic integration is stable and hereditary. Corresponding methods are described, (Jenes 1993; Potrykus 1991).

## PREFERRED GENETIC COMPONENTS

45 Preferably, the genetic component (e.g., the T-DNA) inserted into the genome of the target plant comprises at least one expression cassette, which may – for example – facilitate expression of selection markers, trait genes, antisense RNA or double-stranded RNA. Preferably said expression cassettes comprise a promoter sequence functional in plant cells operatively linked to a nucleic acid sequence which – upon ex-

pression – confers an advantageous phenotype to the so transformed plant. The person skilled in the art is aware of numerous sequences which may be utilized in this context, e.g. to increase quality of food and feed, to produce chemicals, fine chemicals or pharmaceuticals (e.g., vitamins, oils, carbohydrates; Dunwell 2000), conferring resistance to herbicides, or conferring male sterility. Furthermore, growth, yield, and resistance against abiotic and biotic stress factors (like e.g., fungi, viruses or insects) may be enhanced. Advantageous properties may be conferred either by overexpressing proteins or by decreasing expression of endogenous proteins by e.g., expressing a corresponding antisense (Sheehy 1988; US 4,801,340; Mol 1990) or double-stranded RNA (Matzke 2000; Fire 1998; Waterhouse 1998; WO 99/32619; WO 99/53050; WO 00/68374; WO 00/44914; WO 00/44895; WO 00/49035; WO 00/63364).

For expression in plants, plant-specific promoters are preferred. The term "plant-specific promoter" is understood as meaning, in principle, any promoter which is capable of governing the expression of genes, in particular foreign genes, in plants or plant parts, plant cells, plant tissues or plant cultures. In this context, expression can be, for example, constitutive, inducible or development-dependent. The following are preferred:

a) Constitutive promoters

"Constitutive" promoters refers to those promoters which ensure expression in a large number of, preferably all, tissues over a substantial period of plant development, preferably at all times during plant development. A plant promoter or promoter originating from a plant virus is especially preferably used. The promoter of the CaMV (cauliflower mosaic virus) 35S transcript (Franck 1980; Odell 1985; Shewmaker 1985; Gardner 1986) or the 19S CaMV promoter (US 5,352,605; WO 84/02913; Benfey 1989) are especially preferred. Another suitable constitutive promoter is the rice actin promoter (McElroy 1990), Rubisco small subunit (SSU) promoter (US 4,962,028), the legumin B promoter (GenBank Acc. No. X03677), the promoter of the nopalin synthase from Agrobacterium, the TR dual promoter, the OCS (octopine synthase) promoter from Agrobacterium, the ubiquitin promoter (Holtorf et al. 1995), the ubiquitin 1 promoter (Christensen et al. 1989, 1992; Bruce et al. 1989), the Smas promoter, the cinnamyl alcohol dehydrogenase promoter (US 5,683,439), the promoters of the vacuolar ATPase subunits, the pEMU promoter (Last et al. 1991); the MAS promoter (Velten et al. 1984) and maize H3 histone promoter (Lepetit et al. 1992; Atanassova et al. 1992), the promoter of the Arabidopsis thaliana nitrilase-1 gene (GenBank Acc. No.: U38846, nucleotides 3862 to 5325 or else 5342) or the promoter of a proline-rich protein from wheat (WO 91/13991), and further promoters of genes whose constitutive expression in plants, especially in monocot or Gramineae plants is known to the skilled worker. The maize ubiquitin promoter is particularly preferred in wheat and barley.

b) Tissue-specific or tissue-preferred promoters

Furthermore preferred are promoters with specificities for seeds, such as, for example, the phaseolin promoter (US 5,504,200; Bustos et al. 1989, Murai et al. 1983; Sengupta-Gopalan et al. 1985), the promoter of the 2S albumin gene (Joseffson et al. 1987), the legumine promoter (Shirsat 1989), the USP (unknown seed protein) promoter (Bäumlein 1991a), the napin gene promoter (US 5,608,152; Stalberg et al. 1996), the promoter of the sucrose binding proteins (WO 00/26388) or the legumin B4 promoter (LeB4; Bäumlein et al. 1991b, 1992), the Arabidopsis oleosin promoter (WO 98/45461),

and the Brassica Bce4 promoter (WO 91/13980). Promoters which are furthermore preferred are those which permit a seed-specific expression in monocots such as maize, barley, wheat, rye, rice and the like. The promoter of the *lpt2* or *lpt1* gene (WO 95/15389, WO 95/23230) or the promoters described in WO 99/16890 (promoters of the hordein gene, the glutelin gene, the oryzin gene, the prolamin gene, the gliadin gene, the glutelin gene, the zein gene, the casirin gene or the secalin gene) can advantageously be employed. Further preferred are a leaf-specific and light-induced promoter such as that from *cab* or *rubisco* (Simpson 1985; Timko 1985); an anther-specific promoter such as that from *LAT52* (Twell et al. 1989b); a pollen-specific promoter such as that from *Zm13* (Guerrero et al. 1993); and a microspore-preferred promoter such as that from *apg* (Twell et al. 1993).

c) Chemically inducible promoters

The expression cassettes may also contain a chemically inducible promoter (review article: Gatz et al. 1997), by means of which the expression of the exogenous gene in the plant can be controlled at a particular point in time. Such promoters such as, for example, the *PRP1* promoter (Ward et al. 1993), a salicylic acid-inducible promoter (WO 95/19443), a benzenesulfonamide-inducible promoter (EP 0 388 186), a tetracyclin-inducible promoter (Gatz et al. 1991, 1992), an abscisic acid-inducible promoter EP 0 335 528) or an ethanol-cyclohexanone-inducible promoter (WO 93/21334) can likewise be used. Also suitable is the promoter of the glutathione-S transferase isoform II gene (*GST-II-27*), which can be activated by exogenously applied safeners such as, for example, *N,N*-diallyl-2,2-dichloroacetamide (WO 93/01294) and which is operable in a large number of tissues of both monocots and dicots. Further exemplary inducible promoters that can be utilized in the instant invention include that from the *ACE1* system which responds to copper (Mett et al. 1993); or the *ln2* promoter from maize which responds to benzenesulfonamide herbicide safeners (Hershey et al. 1991; Gatz et al. 1994). A promoter that responds to an inducing agent to which plants do not normally respond can be utilized. An exemplary inducible promoter is the inducible promoter from a steroid hormone gene, the transcriptional activity of which is induced by a glucocorticosteroid hormone (Scheda et al. 1991).

Particularly preferred are constitutive promoters. Furthermore, further promoters may be linked operably to the nucleic acid sequence to be expressed, which promoters make possible the expression in further plant tissues or in other organisms, such as, for example, *E. coli* bacteria. Suitable plant promoters are, in principle, all of the above-described promoters.

The genetic component and/or the expression cassette may comprise further genetic control sequences in addition to a promoter. The term "genetic control sequences" is to be understood in the broad sense and refers to all those sequences which have an effect on the materialization or the function of the expression cassette according to the invention. For example, genetic control sequences modify the transcription and translation in prokaryotic or eukaryotic organisms. Preferably, the expression cassettes according to the invention encompass a promoter functional in plants 5'-upstream of the nucleic acid sequence in question to be expressed recombinantly, and 3'-downstream a terminator sequence as additional genetic control sequence and, if appropriate, further customary regulatory elements, in each case linked operably to the nucleic acid sequence to be expressed recombinantly.

Genetic control sequences furthermore also encompass the 5'-untranslated regions, introns or noncoding 3'-region of genes, such as, for example, the actin-1 intron, or the Adh1-S introns 1, 2 and 6 (general reference: The Maize Handbook, Chapter 116, Freeling and Walbot, Eds., Springer, New York (1994)). It has been demonstrated that they may play a significant role in the regulation of gene expression. Thus, it has been demonstrated that 5'-untranslated sequences can enhance the transient expression of heterologous genes. Examples of translation enhancers which may be mentioned are the tobacco mosaic virus 5' leader sequence (Gallie et al. 1987) and the like. Furthermore, they may promote tissue specificity (Rouster et al. 1998).

The expression cassette may advantageously comprise one or more enhancer sequences, linked operably to the promoter, which make possible an increased recombinant expression of the nucleic acid sequence. Additional advantageous sequences, such as further regulatory elements or terminators, may also be inserted at the 3' end of the nucleic acid sequences to be expressed recombinantly. Polyadenylation signals which are suitable as control sequences are plant polyadenylation signals, preferably those which essentially correspond to T-DNA polyadenylation signals from *Agrobacterium tumefaciens*, in particular the OCS (octopin synthase) terminator and the NOS (nopal synthase) terminator.

Control sequences are furthermore to be understood as those permitting removal of the inserted sequences from the genome. Methods based on the cre/lox (Sauer B. 1998; Odell 1990; Dale 1991), FLP/FRT (Lysnik 1993), or Ac/Ds system (Wader et al. 1987; US 5,225,341; Baker et al. 1987; Lawson et al. 1994) permit a - if appropriate tissue-specific and/or inducible - removal of a specific DNA sequence from the genome of the host organism. Control sequences may in this context mean the specific flanking sequences (e.g., lox sequences), which later allow removal (e.g., by means of cre recombinase).

The genetic component and/or expression cassette of the invention may comprise further functional elements. The term functional element is to be understood in the broad sense and refers to all those elements which have an effect on the generation, amplification or function of the genetic component, expression cassettes or recombinant organisms according to the invention. Functional elements may include for example (but shall not be limited to):

#### 1. Selection markers

Selection markers are useful to select and separate successfully transformed or homologous recombined cells.

##### 1.1 Positive selection markers

Selection markers confer a resistance to a biocidal compound such as a metabolic inhibitor (e.g., 2-deoxyglucose-6-phosphate, WO 98/45456), antibiotics (e.g., kanamycin, G 418, bleomycin or hygromycin) or herbicides (e.g., phosphinothricin or glyphosate). Especially preferred selection markers are those which confer resistance to herbicides. Examples which may be mentioned are:

- Phosphinothricin acetyltransferases (PAT; also named Bialophos<sup>®</sup> resistance; bar; de Block 1987; . EP 0 333 033; US 4,975,374)
  - 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) conferring resistance to Glyphosate<sup>®</sup> (N-(phosphonomethyl)glycine) (Shah 1986)
  - Glyphosate<sup>®</sup> degrading enzymes (Glyphosate<sup>®</sup> oxidoreductase; gox),
  - Dalapon<sup>®</sup> inactivating dehalogenases (deh)
  - sulfonylurea- and imidazolinone-inactivating acetolactate synthases (for example mutated ALS variants with, for example, the S4 and/or Hra mutation)
  - Bromoxynil<sup>®</sup> degrading nitrilases (bxn)
  - Kanamycin- or. G418- resistance genes (NPTII; NPTI) coding e.g., for neomycin phosphotransferases (Fraley 1983)
  - 2-Desoxyglucose-6-phosphate phosphatase (DOG<sup>R</sup>1-Gene product; WO 98/45456; EP 0 807 836) conferring resistance against 2-desoxyglucose (Randez-Gil 1995).
  - hygromycin phosphotransferase (HPT), which mediates resistance to hygromycin (Vanden Elzen 1985).
  - dihydrofolate reductase (Eichholtz 1987)
- Additional positive selectable marker genes of bacterial origin that confer resistance to antibiotics include the *aadA* gene, which confers resistance to the antibiotic spectinomycin, gentamycin acetyl transferase, streptomycin phosphotransferase (SPT), aminoglycoside-3-adenyl transferase and the bleomycin resistance determinant (Hayford 1988; Jones 1987; Svab 1990; Hille 1986).
- Especially preferred are selection markers that confer resistance against the toxic effects imposed by D-amino acids like e.g., D-alanine and D-serine (WO 03/060133). Especially preferred as negative selection marker in this contest are the *daa1* gene (EC: 1.4. 3.3 : GenBank Acc.-No.: U60066) from the yeast *Rhodotorula gracilis* (*Rhodospiridium toruloides*) and the *E. coli* gene *dsdA* (D-serine dehydratase (D-serine deaminase) [EC: 4.3. 1.18; GenBank Acc.-No.: J01603).
- Genes like isopentenyltransferase from *Agrobacterium tumefaciens* (strain:PO22; Genbank Acc.-No.: AB025109) may – as a key enzyme of the cytokinin biosynthesis – facilitate regeneration of transformed plants (e.g., by selection on cytokinin-free medium). Corresponding selection methods are described (Ebinuma H et al. (2000) Proc Natl Acad Sci USA 94:2117-2121; Ebinuma H et al. (2000) Selection of Marker-free transgenic plants using the oncogenes (*ipt*, *rol A*, B, C) of *Agrobacterium* as selectable markers, In Molecular Biology of Woody Plants. Kluwer Academic Publishers). Additional positive selection markers, which confer a growth advantage to a transformed plant in comparison with a non-transformed one, are described e.g., in EP-A 0 601 092. Growth stimulation selection markers may include (but shall not be limited to)  $\beta$ -Glucuronidase (in combination with e.g., a cytokinin glucuronide), mannose-6-phosphate isomerase (in combination with mannose), UDP-galactose-4-epimerase (in combination with e.g., galactose), wherein mannose-6-phosphate isomerase in combination with mannose is especially preferred.

## 1.2) Negative selection marker

Negative selection markers are especially suitable to select organisms with defined deleted sequences comprising said marker (Koprek et al. 1999). Examples for negative selection marker comprise thymidin kinases (TK), cytosine deaminases (Gleave et al. 1999; Perera et al. 1993; Stougaard 1993), cytochrom P450 proteins (Koprek et al. 1999), haloalkane dehalogenases (Naested 1999), iaaH gene products (Sundaresan et al. 1995), cytosine deaminase codA (Schlaman & Hooykaas 1997), or tms2 gene products (Fedoroff & Smith 1993).

### 2) Reporter genes

Reporter genes encode readily quantifiable proteins and, via their color or enzyme activity, make possible an assessment of the transformation efficacy, the site of expression or the time of expression. Very especially preferred in this context are genes encoding reporter proteins (Schenborn 1999) such as the green fluorescent protein (GFP) (Sheen 1995; Haseloff 1997; Reichel 1996; Tian 1997; WO 97/41228; Chui 1996; Lefel 1997), chloramphenicol transferase, a luciferase (Ow 1986; Millar 1992), the aequorin gene (Prasher 1985),  $\beta$ -galactosidase, R locus gene (encoding a protein which regulates the production of anthocyanin pigments (red coloring) in plant tissue and thus makes possible the direct analysis of the promoter activity without addition of further auxiliary substances or chromogenic substrates (Dellaporta 1988; Ludwig 1990), with  $\beta$ -glucuronidase (GUS) being very especially preferred (Jefferson 1987a,b).  $\beta$ -glucuronidase (GUS) expression is detected by a blue color on incubation of the tissue with 5-bromo-4-chloro-3-indolyl- $\beta$ -D-glucuronic acid, bacterial luciferase (LUX) expression is detected by light emission; firefly luciferase (LUC) expression is detected by light emission after incubation with luciferin; and galactosidase expression is detected by a bright blue color after the tissue is stained with 5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside. Reporter genes may also be used as scorable markers as alternatives to antibiotic resistance markers. Such markers are used to detect the presence or to measure the level of expression of the transferred gene. The use of scorable markers in plants to identify or tag genetically modified cells works well only when efficiency of modification of the cell is high.

3) Origins of replication, which ensure amplification of the expression cassettes or vectors according to the invention in, for example, *E. coli*. Examples which may be mentioned are ORI (origin of DNA replication), the pBR322 ori or the P15A ori (Maniatis 1989).

4) Elements which are necessary for *Agrobacterium*-mediated plant transformation, such as, for example, the right or left border of the T-DNA or the vir region.

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the composition, methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described



herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference for all purposes.

#### Sequences

1. SEQ ID NO: 1 Binary vector pUGAB7
2. SEQ ID NO: 2 Binary vector pSUNUbi-GFP
3. SEQ ID NO: 3 Binary vector pSUNUbiGUS

#### Examples:

##### General methods:

Unless otherwise specified, all chemicals were from Fluka (Buchs), Merck (Darmstadt), Roth (Karlsruhe), Serva (Heidelberg) and Sigma (Deisenhofen). Restriction enzymes, DNA-modifying enzymes and molecular biological kits were from Amersham-Pharmacia (Freiburg), Biometra (Göttingen), Roche (Mannheim), New England Biolabs (Schwalbach), Novagen (Madison, Wisconsin, USA), Perkin-Elmer (Weiterstadt), Qiagen (Hilden), Stratagen (Amsterdam, Netherlands), Invitrogen (Karlsruhe) and Ambion (Cambridgeshire, United Kingdom). The reagents used were employed in accordance with the manufacturer's instructions.

For example, oligonucleotides can be synthesized chemically in the known manner using the phosphoramidite method (Voet, 2nd edition, Wiley Press New York, pages 896-897). The cloning steps carried out for the purposes of the present invention such as, for example, restriction cleavages, agarose gel electrophoreses, purification of DNA fragments, transfer of nucleic acids to nitrocellulose and nylon membranes, linking DNA fragments, transformation of *E. coli* cells, bacterial cultures, multiplication of phages and sequence analysis of recombinant DNA, are carried out as described (Maniatis 1989). Recombinant DNA molecules were sequenced using an ABI laser fluorescence DNA sequencer following the method of Sanger (Sanger 1977).

Plant material: Culture of wheat (*Triticum aestivum* L.) plants, emasculation of spikes and hand-pollination were conducted as described (Kumlehn et al. 1997). As genotypes, the German winter-type cultivar Certo and the spring-type Bobwhite were used.

DNA-Constructs: The *Agrobacterium* binary constructs pSUNUbiGUS (SEQ ID NO: 3) and pSUNUbiGFP (SEQ ID NO: 2) for marker gene expression were generated using the pSUN300, a derivative of the pSUN binary vector (WO 02/00900). pSUNUbiGUS and pSUNUbiGFP were generated by insertion of a Ubi-1 promoter/ GUS(intron)/ 35S-T expression cassette and a Ubi-1 promoter/ GFP/ 35S-T expression cassette respectively into the binary vector. Thereafter the P-nos/ nptII/ nosT cassette of pSUN300 was replaced by a Ubi-1-promoter/pat/35S-T cassette to apply both binary plasmids to wheat transformation.

### Example 1: Isolation of wheat zygotes

#### 1.1. Donor plant material

Plants of different genotypes of spring (Bobwhite, Canon, Salmon) or winter (Certo) wheat (*Triticum aestivum* L.) were grown at 12/ 10 °C day/ night temperature with 10 hours of daily illumination with 30,000 lx in climatic chambers followed by 18/ 14 °C day/night temperature and 16 hours daily illumination with a minimum of 25,000 lx in green houses. It was necessary to vernalize the seedlings of cv. Certo for 6 to 8 weeks at 4 °C and short day illumination of 2,000 lx prior to transfer to said growth conditions necessary for further development.

#### 1.2. Emasculation

Emasculation was performed when spikes had emerged from the flag leaf after approximately 10 to 12 weeks after seed sowing. First of all, the underdeveloped lowest and uppermost spikelets as well as the central florets were removed from the wheat spikes. Then the outer glumes and finally the anthers from the remaining wheat flowers were removed. To prevent free cross pollination, the emasculated spikes were covered with polyethylene bags. Emasculation was performed 2 days prior to the date at which anthesis would be commencing.

#### 1.3. Pollination

Two days after emasculation, the florets were manually pollinated using anthers and pollen freshly harvested from other wheat spikes.

#### 1.4. Surface sterilization of spikes

One to five hours after manual pollination, the spikes were cut for further processing. The spikes were surface sterilized in 2% sodium hypochlorite (v/v) in addition to 0.03% Tween 20 for 10 minutes followed by 5 times washing with sterile distilled water. Excess water was removed from the spikes with the help of sterilized tissue paper.

#### 1.5. Ovule tip preparation

Pistils were dissected from a spike and placed in a sterile glass petri dish. Still attached parts (filaments, lodicules or palea) were carefully removed by use of a standard preparation microscope. The ovary tip was cut off starting from the pistil's adaxial side at a position near the entrance of the vascular bundle. The cut ovary tip was immediately transferred to 0.55 M mannitol solution in a 35-mm petri dish (Greiner, Frickenhausen/Germany). Upon submergence of the ovary tips, the ovule tips comprising the inner integumental layer, nucellus tissue and a part of the embryo sac were dissected and the pericarp together with the outer integument removed from the solution.

#### 1.6. Zygote isolation and transfer

Further processing was performed by use of an inverted microscope. An ovule tip was fixed onto the bottom of the dish with the help of an L-bended fine-tipped glass needle and the zygote pushed out of the ovule tip by means of a second glass needle. Since the wheat embryo sac is not as tightly embedded in nucellus tissue as, for instance, the maize embryo sac, the zygote can be isolated efficiently without use of cell-wall-degrading enzymes. With this method it was possible to routinely recover viable zygotes from about 75% of the excised pistils in genotypes of both winter and spring

types. The isolated zygotes had a diameter of around 75  $\mu\text{m}$ . Upon isolation, the zygotes were distributed to Millicell inserts of 12 mm diameter and 0.4  $\mu\text{m}$  pore size (Millicore, Milshiem/France) placed in 35-mm culture dishes or in chamber slides. Up to 10 zygotes were cultured per Millicell insert. Uptake and release of the zygotes were conducted by means of a glass capillary connected to a manually controlled Cell Tram (Eppendorf, Germany) and filled with 0.55 M mannitol solution.

#### Example 2 : Barley immature pollen culture

Barley tillers of cv. Igri were harvested when the tips of the awns were just emerging from the sheath of the flag leaf. The tillers were surface-sterilized by spraying with 70% ethanol. After having released a spike from the flag leaf, the awns were removed with forceps and five spikes were put onto a 70-mm sterile filter paper disc which had been dipped once in sterile distilled water and placed into a 9-cm petri dish. The dishes were sealed and incubated at 4°C for about 4 weeks in the dark. To isolate immature pollen, 10 to 15 pretreated spikes were chopped down to 1 to 2 cm pieces to be transferred to a pre-cooled (4°C) Waring Blendor. The Blendor was filled with 20 ml of ice-cold 0.4 M mannitol and the spikes blended 2 times for 5 seconds at low speed. The solution was poured through a 100- $\mu\text{m}$  nylon mesh into an autoclaved Magenta box. The debris was slightly squashed, rinsed with 5 ml of 0.4 M mannitol and transferred back into the blendor. Blending was repeated using 10 ml of 0.4 mannitol solution. The interior of the blendor was rinsed with 5 ml of mannitol solution which was subsequently poured through the remaining debris in the mesh. The pollen suspension was transferred from the Magenta box into a 50-ml polypropylene tube which was closed with a screw cap and centrifuged at 100 g for 10 min at 4°C. The supernatant was removed and the pellet resuspended in 5 ml of 0.55 M maltose. The suspension was equally distributed to two 12-ml centrifuge tubes and cautiously overlaid with 2 x 1.5 ml of 0.4 M mannitol. After gradient centrifugation at 100 g for 10 minutes, the interphase bands containing the vacuolated pollen were transferred to a 50-ml polypropylene tube which were filled up to 20 ml with 0.4 M mannitol. The population density was assessed by means of a haemocytometer and the vacuolated pollen centrifuged again at 100 g for 10 minutes. The supernatant was discarded and the population density was adjusted to  $2.5 \times 10^5$  per ml by adding KBP medium (Table 1). One-ml aliquots of the suspension were distributed to 35-mm petri dishes. The cultures were incubated at 25°C in the dark.

#### Example 3: Growth of *Agrobacterium tumefaciens*

A cryostock of LBA4404/ pSB1/ pUGAB7 consisting of 200  $\mu\text{l}$  of a growing *Agrobacterium* culture with a density of  $10^9$  cfu per ml and 200  $\mu\text{l}$  of 15 % sterile glycerol solution was taken from -80°C and given into 10 ml of CPY medium supplemented with 10 mg/l tetracyclin and 150 mg /l spectinomycin. The culture was incubated at 28°C on a rotary shaker at 200 rpm in the dark for about 24 hours. The *Agrobacteria* were pelleted at 4000 rpm for 10 minutes and the supernatant removed. The pellet was resuspended in fresh medium without antibiotics and incubated on a rotary shaker ( ~60 rpm) at 25°C in the dark for 1 to 3 hours.

**Example 4: Co-culture of isolated wheat zygotes with *Agrobacterium tumefaciens* and early embryonic development**

Two Millicell inserts were placed in a 35-mm petri dish with 1.5 ml of ZT1 medium (Table 1). One ml of ZT1 medium containing about 5000 sporophytically developing microcalli taken from a 2-week old barley pollen culture were added. The Millicell inserts were filled with 200  $\mu$ l of ZT1 medium each. Upon isolation in 0.55 M mannitol, five to 10 wheat zygotes of cvs. Certo or Bobwhite were transferred in about 100 nl of 0.55 M mannitol solution by means of a glass capillary interfaced by teflon tube to a Cell Tram (Eppendorf, Hamburg/Germany) into one of the Millicell inserts. To ensure sufficient space for embryo formation, not more than ten zygotes were cultivated per insert.

A culture of *Agrobacterium tumefaciens* strain LBA4404/ pSB1 (WO 95/06722) carrying the binary vector pUGAB7 (Fig. 2) or pYF133 (Fang et al. 2002) was adjusted to a population density of  $10^6$  cfu per ml co-culture medium. Two  $\mu$ l of the resulting suspension was transferred to the Millicell insert that contained the isolated zygotes to give a final *Agrobacterium* density of  $10^4$  cfu per ml within the Millicell insert. The co-culture dishes were incubated on a rotary shaker (~60 rpm) at 21°C in the dark for 48 hours. After this co-culture period, the zygotes were transferred by means of the above mentioned capillary system in about 100 nl of medium to the second Millicell insert of the same petri dish. The Millicell insert which had been used for co-culture was removed. Upon transfer of the zygotes or zygote-derived structures, the ZT1-medium inside the petri dish was supplemented with 13.5  $\mu$ l of a Timentin (Duchefa, Netherlands) stock solution (40 mg/ml) and 18.9  $\mu$ l of a 1 M  $\text{CaCl}_2$  stock solution in order to stop the development of the *Agrobacteria*. Furtheron, 32.4  $\mu$ l of a 0.25 M glutamine solution was added to the medium to support embryonic development.

Alternatively, the zygotes were co-cultured in Millicell inserts using ZT2 medium (Tab.1). Fifty percent of this medium had been conditioned with immature wheat pistils dissected at the late microspore stage of the florets. To this end, ten pistils were cultured per ml at 26 °C for 1 to 2 weeks in the dark. For co-culture, five precultured pistils per ml were placed into the medium outside the Millicell insert. The other co-culture conditions were as described above. After co-culture, Timentin was added to the medium to obtain a final concentration of 150 mg/l to prevent further surviving of the *Agrobacteria*. As a result of this procedure, transgene expression (gfp under control of the maize Ubiquitin 1 promoter) was observed in two out of 20 zygotes immediately after co-culture (Fig. 5).

**Example 5: Regeneration and analysis of transgenic plants from isolated wheat zygotes co-cultivated with *Agrobacteria***

About 4 weeks after co-culture, zygote-derived embryos of at least 1 mm in size were transferred on to K4NT medium (Table 1) for regeneration. Genomic DNA from regenerated plantlets was isolated and analysed for presence and integration of the transgenes by PCR and Southern blot, respectively.

Table 1

Components [mg·l <sup>-1</sup> ]	N6Z	KBP	ZT1	ZT2	K4NT
NH <sub>4</sub> NO <sub>3</sub>	-	80	-	80	320
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	231	-	231	-	-
KNO <sub>3</sub>	1,415	2,022	1,415	2,022	3,640
KH <sub>2</sub> PO <sub>4</sub>	200	340	1,360	340	340
CaCl <sub>2</sub> ·2H <sub>2</sub> O	83	441	441	441	441
MgSO <sub>4</sub> ·7H <sub>2</sub> O	93	246	93	246	246
NaFeEDTA	25	27.5	25	27.5	27.5
MnSO <sub>4</sub> ·4H <sub>2</sub> O	4.0	11.2	4.0	11.2	11.2
H <sub>3</sub> BO <sub>3</sub>	0.5	3.1	0.5	3.1	3.1
ZnSO <sub>4</sub> ·7H <sub>2</sub> O	0.5	7.2	0.5	7.2	7.2
Na <sub>2</sub> MoO <sub>4</sub> ·2H <sub>2</sub> O	0.025	0.12	0.025	0.12	0.12
CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.025	0.025	0.025	0.025	1.25
CoCl <sub>2</sub> ·6H <sub>2</sub> O	0.025	0.024	0.025	0.024	0.024
KI	-	0.17	-	0.17	0.17
retinol	0.01	0.01	0.01	0.01	-
thiamine·HCl	1.0	1.0	1.0	1.0	10.0
nicotinic acid	1.0	1.0	1.0	1.0	1.0
riboflavin	0.2	0.2	0.2	0.2	-
Ca-pantothenate	1.0	1.0	1.0	1.0	-
folic acid	0.4	0.4	0.4	0.4	-
pyridoxine·HCl	1.0	1.0	1.0	1.0	1.0
cobalamine	0.02	0.02	0.02	0.02	-
ascorbic acid	2.0	2.0	2.0	2.0	-
calciferol	0.01	0.01	0.01	0.01	-
biotin	0.01	0.01	0.01	0.01	-
cholin chloride	1.0	1.0	1.0	1.0	-
p-aminobenzoic acid	0.02	0.02	0.02	0.02	-
myo-inositol	100	100	100	100	-
malic acid	400	-	-	-	-
citric acid	40	-	-	-	-
fumaric acid	40	-	-	-	-
Na-pyruvate	20	-	-	-	-
glutamine	1,000	438.4	438.4	146	146
casein hydrolysate	250	-	-	-	-
myo-inositol	100	-	100	100	100
xylose	150	-	150	-	-
glucose	85,000	-	90,000	85,000	-
maltose·H <sub>2</sub> O	-	90,000	-	-	36,000
2,4-D	0.2	-	-	0.2	-
BAP	-	0.22	-	-	0.22
MES	-	-	1,950	1,950	-
Acetosyringone	-	-	100	100	-
Timentin™	-	-	-	-	150
Phytigel	-	-	-	-	4,000
pH	5.7	5.8	5.9	5.9	5.8

## REFERENCES

The references listed below and all references cited herein are incorporated herein by reference to the extent that they supplement, explain, provide a background for, or teach methodology, techniques, and/or compositions employed herein.

1. Allington "Micromanipulation of the Unfixed Cereal Embryo Sac," in THE EXPERIMENTAL MANIPULATION OF OVULE TISSUES 39-51 (Longman, New York, 1985)
2. An et al. (1985) EMBO J 4:277-287
- 10 3. Ashby et al. (1988) J. Bacteriol. 170: 4181-4187
4. Atanassova et al. (1992) Plant J 2(3): 291-300
5. Ausubel FM et al. (1987) Current Protocols in Molecular Biology, Greene Publishing Assoc. and Wiley Interscience
6. Baker et al. (1987) EMBO J 6: 1547-1554
- 15 7. Bäumlein et al. (1992) Plant J 2(2):233-9;
8. Bäumlein H et al. (1991a) Mol Gen Genet 225(3):459-467
9. Bäumlein H et al. (1991b) Mol Gen Genet 225:121-128
10. Becker et al. (1992) Plant Mol. Biol. 20: 49
11. Becker et al. (1994) Plant J., 5:299-307,
- 20 12. Benfey et al.(1989) EMBO J 8:2195-2202
13. Bevan et al. (1984) Nucl Acid Res 12,8711-8720
14. Bolton et al. (1986) Science 232: 983-985;
15. Bruce et al. (1989) Proc Natl Acad Sci USA 86:9692-9696
16. Bustos MM et al. (1989) Plant Cell 1(9):839-53)
- 25 17. Chen and Winans (1991) J. Bacteriol. 173: 1139-1144
18. Christensen et al. (1989) Plant Mol. Biol. 12: 619-632
19. Christensen et al. (1992) Plant Mol Biol 18:675-689
20. Christou et al. (1988) Plant Physiol 87:671-674
21. Christou et al. (1991) Bio/Technology 9:957-962
- 30 22. Chu CC et al. (1990) Plant Sci 66:255- 767
23. Chui WL et al. (1996) Curr Biol 6:325-330;
24. Creissen et al. (1991) Plant J. 2: 129
25. Dale & Ow (1991) Proc Nat'l Acad Sci USA 88:10558-10562
26. Datta et al. (1990b) Bio/Technology 8:736-740
- 35 27. Datta SK & Wenzel G (1987) Plant Sci 48:49-54
28. Datta SK et al. (1990a) Plant Sci. 67, 83-88
29. Davey et al. (1991) J Exp Bot 42:1129-1169
30. de Block et al. (1987) EMBO J 6:2513-2518
31. De la Pena et al. (1987) Nature 325:274-276
- 40 32. Deblaere et al. (1985) Nucl Acids Res 13:4777-4788
33. Dekeyser et al. (1989) Plant Physiol 90:217-223
34. Della-Cioppa et al. (1987) Bio/Technology 5:579-584
35. Dellaporta et al. (1988) In: Chromosome Structure and Function: Impact of New Concepts, 18th Stadler Genetics Symposium, 11:263-282
- 45 36. Du et al. (1989) Genet Manip Plants 5:8-12
37. Dunwell JM (2000) J Exp Bot 51 Spec No:487-96
38. Eichholtz et al. (1987) Somatic Cell and Molecular Genetics 13, 67-76
39. Engell K (1989). Nord. J. Bot. 9, 265-280.
40. EP-A1 0 120 516

41. EP-A1 0 120 516
42. EP-A1 0 333 033
43. EP-A1 0 335 528
44. EP-A1 0 388 186
- 5 45. EP-A1 0 807 836
46. Evans DA & Bravo JE (1983). Protoplast isolation and culture. In: D.A. Evans, W.R. Sharp, Ammirato and Y. Yamada eds. Handbook of Plant Cell Culture. Macmillan Publishing Company, V. 1, 124-176.
47. Fang Y et al. (2002) J. Plant Physiol. 159 1131-1138
- 10 48. Fedoroff NV & Smith DL (1993) Plant J 3:273- 289
49. Fire A. et al (1998) Nature 391:806-811
50. Fontes et al. (1991) Plant Cell 3:483-496
51. Fraley et al. (1983) Proc Natl Acad Sci USA 80:4803
52. Franck et al. (1980) Cell 21:285-294
- 15 53. Fromm et al. (1986) Nature 319:791-793
54. Fromm et al. (1990) Bio/Technology 8:833-839
55. Gallie et al. (1987) Nucl Acids Res 15:8693-8711
56. Gardner et al. (1986) Plant Mol Biol 6:221-228
57. Gatz et al. (1991) Mol Gen Genetics 227:229-237
- 20 58. Gatz et al. (1992) Plant J 2:397-404
59. Gatz et al. (1994) Mol Gen Genetics 243:32-38
60. Gatz et al. (1997) Annu Rev Plant Physiol Plant Mol Biol 48:89-108
61. Gelvin et al. (Eds) (1990) Plant Molecular Biology Manual; Kluwer Academic Publisher, Dordrecht, The Netherlands
- 25 62. GenBank Acc. No. X03677
63. GenBank Acc. No.: U38846, nucleotides 3862 to 5325 or else 5342
64. Gleave AP et al. (1999) Plant Mol Biol. 40(2):223-35
65. Gordon-Kamm et al. (1990) Plant Cell 2:603-618
66. Gould et al. (1989) J Cell Biol 108:1657
- 30 67. Gould et al. (1991) Plant Physiol 95(2):426-434
68. Gruber et al. (1993) "Vectors for Plant Transformation," in METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY; pp.89-119.
69. Guerrero et al. (1993) Mol Gen Genet 224:161-168
70. Guivarc'h et al. (1993) Protoplasma 174: 10-18
- 35 71. Hajdukiewicz et al. (1994) Plant Mol Biol 25:989-994
72. Hansen et al. (1994) Proc. Natl. Acad. Sci. USA 91:7603-7607
73. Haseloff et al. (1997) Proc Natl Acad Sci USA 94(6):2122-2127;
74. Hayakawa et al. (1992) Proc Natl Acad Sci USA 89:9865-9869
75. Hayford et al. (1988) Plant Physiol. 86:1216
- 40 76. Hershey et al. (1991) Mol Gen Genetics 227:229-237
77. Hiei et al. (1994) Plant J 6: 271-282
78. Hille et al. (1986) Plant Mol. Biol. 7:171
79. Hoekema (1985) In: The Binary Plant Vector System, Offsetdrukkerij Kanters B.V., Alblisserdam, Chapter V
- 45 80. Hoekema et al. (1983) Nature 303:179-181
81. Holm PB et al. (1994) Plant Cell 6: 531-543
82. Holsters et al. (1978) Mol Gen Genet 163:181-187
83. Holtorf S et al. (1995) Plant Mol Biol 29:637-649
84. Hood et al. (1986) J Bacteriol 168:1291-1301



85. Horsch RB et al. (1985) *Science* 225: 1229f
86. Ishida Y et al. (1996) *Nature Biotech* 745-750
87. Jacq et al. (1993) *Plant Cell Reports* 12: 621-624
88. James et al. (1993) *Plant Cell Reports* 12: 559-563
- 5 89. Jarchow et al. (1991), *Proc. Natl. Acad. Sci. USA* 88:10426-10430
90. Jefferson (1987b) *Plant Mol. Bio. Rep.*, 5:387-405
91. Jefferson et al. (1987a) *EMBO J.*, 6:3901-3907
92. Jenes B et al. (1993) *Techniques for Gene Transfer*, in: *Recombinant Plants*, Vol. 1, Engineering and Utilization, edited by SD Kung and R Wu, Academic Press, pp. 128-143
- 10 93. Johri BM et al. (1992) *Comparative embryology of angiosperms*. Springer, Berlin
94. Jones et al. (1987) *Mol. Gen. Genet.*, 210:86
95. Joseffson LG et al. (1987) *J Biol Chem* 262:12196-12201
96. Kado (1991) *Crit Rev Plant Sci* 10:1
97. Kao KN & Michayluk MR (1975) *Planta (Berl.)* 126, 105-110.
- 15 98. Kasha et al. (1990) *Gene Manip. Plant Improv. II*, 213-239
99. Klapwijk et al. (1980) *J. Bacteriol.*, 141,128-136
100. Knox et al. (1987) *Plant Mol. Biol.* 9:3-17
101. Köhler and Wenzel (1978) *J. Plant Physiol.* 121: 181-191
102. Koncz & Schell (1986) *Mol Gen Genet* 204:383-396
- 20 103. Koop HU, Schweiger HG (1995) *J Plant Physiol* 111: 235-757
104. Koprek T et al. (1999) *Plant J* 19(6): 719-726
105. Kranz E et al. (1991) *Ses Plant Reprod* 4: 12-16
106. Kranz E, Lörz H (1993) *Plant Cell* 5: 739-746
107. Kranz E, Lörz H (1994) *Zygote* 2: 175-13s
- 25 108. Kumlehn et al. (1997) *Plant J.* 12: 1473-1479
109. Kumlehn J et al. (1998) *Planta* 205:327-333
110. Last DI et al. (1991) *Theor. Appl. Genet.* 81, 581-588.
111. Lawson et al. (1994) *Mol Gen Genet* 245:608-615
112. Leduc et al., *Sex Plant Reprod* 8:313-317 (1995)
- 30 113. Leduc N et al. (1996) *Dev Biol* 177: 190-203
114. Leffel SM et al. (1997) *Biotechniques.* 23(5):912-8
115. Lepetit et al., *Mol. Gen. Genet.* 231: 276-285 (1992)
116. Lerner et al., *Plant Physiol.* 91: 124-129 (1989)
117. Li et al. (1992) *Plant Mol Biol* 20:1037-1048
- 35 118. Linsmaier EM and Skoog F (1965) *Physiol. Plant.* 18, 100-127.
119. Ludwig et al. (1990) *Science* 247:449
120. Luo and Wu, *PlantMol. Biol. Rep.*, 6:165-174, 1988.
121. Lysnik et al. (1993) *NAR* 21:969-975
122. Maniatis T, Fritsch EF and Sambrook J (1989) *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor (NY)
- 40 123. Matsuoka et al., *Proc. Nat'l Acad. Sci.* 88: 834 (1991)
124. Matzke MA et al. (2000) *Plant Mol Biol* 43:401-415
125. McElroy et al., *Plant Cell* 2: 163171 (1990)
126. Meinke DW (1993) *Plant Cell* 3: 857-866
- 45 127. Mejza et al. (1993) *Plant Cell Reports* 12: 149-153
128. Mett et al. *PNAS* 90: 4567-4571 (1993)
129. Miki et al. (1993) "Procedures for Introducing Foreign DNA into Plants" in *METHODS IN PLANT MOLECULAR BIOLOGY AND BIOTECHNOLOGY*; pp.67-88
130. Millar et al. (1992) *Plant Mol Biol Rep* 10:324-414

131. Mogensen HL (1982) *Carlsberg Res. Commun.* 47, 313-354.
132. Mol JN et al. (1990) *FEBS Lett* 268(2):427-430
133. Moloney et al., *Plant Cell Reports* 8: 238 (1989)
134. Mooney and Doodwin, *Plant Cell Tissue Organ Culture*, 25:209-218, 1991.
- 5 135. Mordhorst AP, Lorz H (1993) *J Plant Physiol* 131: 485-491
136. Mozo & Hooykaas, *Plant Mol. Biol.* 16 (1991), 917-918.
137. Murai et al., *Science* 23: 476-482 (1983)
138. Murashige T and Skoog F (1962) *Physiol. Plant.* 15, 472-497. (472/473)
139. Naested H (1999) *Plant J* 18:571-576
- 10 140. Nehra et al., *Plant J.*, 5:285-297, 1994.
141. Odell et al. (1985) *Nature* 313:810-812
142. Odell et al. (1990) *Mol Gen Genet* 223:369-378
143. Olhoft PM et al. (2001) *Plant Cell Rep* 20: 706-711
144. Ow et al. (1986) *Science* 234:856-859;
- 15 145. Paszkowski et al. (1984) *EMBO J* 3:2717-2722
146. Perera RJ et al. (1993) *Plant Mol. Biol* 23(4): 793-799;
147. Perl A et al. (1996) *Nature Biotechnol* 14: 624-628
148. Potrykus & Shillito RD (1986) *Methods Enzymol.* 118, 549-578.
149. Potrykus (1990) *Bio/technology.* 8, 535-542.
- 20 150. Potrykus (1991) *Ann Rev Plant Physiol Plant Mol Biol* 42:205-225
151. Prasher et al. (1985) *Biochem Biophys Res Commun* 126(3):1259-1268)
152. Raineri et al. (1990) *Bio/Technology* 8:33
153. Randez-Gil et al. (1995) *Yeast* 11:1233-1240
154. Rasco-Gaunt et al. (2001) *J. Exp. Bot.* 52: 865-874
- 25 155. Reichel et al.(1996) *Proc Natl Acad Sci USA* 93(12):5888-5893;
156. Rhodes CA et al. (1988). *Science* 240, 204-207.
157. Rouster J et al. (1998) *Plant J* 15:435-440
158. Sanford JC (1990) *Physiologia Plantarum* 79:206-209
159. Sanger et al. (1977) *Proc Natl Acad Sci USA* 74:5463-5467
- 30 160. Sauer B (1998) *Methods* 14(4):381-92
161. Sautter et al. (1991) *Bio/Technology*, 9:1080-1085
162. Scheeren-Groot et al. (1994) *J. Bacteriol* 176: 6418-6426
163. Schena et al. (1991) *Proc Nat'l Acad Sci USA* 88:10421
164. Schenborn E, Groskreutz D. (1999) *Mol Biotechnol* 13(1):29-44
- 35 165. Schlaman HRM and Hooykaas PJJ (1997) *Plant J* 11:1377-1385
166. Sengupta-Gopalan et al., *Proc. Nat'l Acad. Sci. USA* 82: 3320-3324 (1985)
167. Shah et al. (1986) *Science* 233: 478
168. Sheehy et al. (1988) *Proc Natl Acad Sci USA* 85: 8805-8809;
169. Sheen et al.(1995) *Plant J* 8(5):777-784;
- 40 170. Shewmaker et al. (1985) *Virology* 140:281-288
171. Shillito et al. (1985) *Bio/Technology*, 3:1099-1103
172. Shimamoto et al. (1989) *Nature* 338:274-276
173. Shirsat A et al. (1989) *Mol Gen Genet* 215(2):326-331
174. Silhavy TJ, Berman ML and Enquist LW (1984) *Experiments with Gene Fusions*, Cold Spring Harbor Laboratory, Cold Spring Harbor (NY)
- 45 175. Simpson et al. (1985) *EMBO J* 4:2723-2729
176. Somers et al. (1992) *Bio/Technology* 10:1589-1594
177. Spencer TM et al. (1990) *Theor Appl Genet* 79, 625-631
178. Stachel et al. (1985) *Nature* 318: 624-629

179. Stalberg K et al. (1996) *Planta* 199:515-519  
180. Stougaard J; (1993) *Plant J* 3:755-761  
181. Svab et al., (1990) *Plant Mol. Biol.* 14:197  
182. The Maize Handbook, Chapter 116, Freeling and Walbot, Eds., Springer, New York  
5 (1994)  
183. Theunis et al. (1991) *Sex Plant Reprod.* 4:145-154  
184. Thomas TL (1993) *Plant Cell* 5:1401-1410  
185. Tian et al. (1997) *Plant Cell Rep* 16:267-271;  
186. Timko et al. (1985) *Nature* 318: 579-582  
10 187. Topfer et al. (1989) *Plant Cell*, 1:133-139  
188. Twell et al. (1983) *Sex. Plant Reprod.* 6: 217-224  
189. Twell et al. (1989a) *Plant Physiol* 91:1270-1274  
190. Twell et al. (1989b) *Mol Gen Genet* 217:240-245  
191. US 4,801,340;  
15 192. US 4,962,028  
193. US 4,975,374)  
194. US 4,940,838  
195. US 5,225,341  
196. US 5,352,605  
20 197. US 5,504,200  
198. US 5,565,350  
199. US 5,608,152;  
200. US 5,683,439  
201. Van Laerebeke et al. (1974) *Nature* 252,169-170  
25 202. Van Wordragen and Dons (1992) *Plant Mol. Biol. Rep.* 10: 12-36  
203. Vanden Elzen et al. (1985) *Plant Mol Biol.* 5:299  
204. Vasil et al. (1992) *Bio/Technology*, 10:667-674  
205. Vasil et al. (1993) *Bio/Technology*, 11:1153-1158  
206. Velten J et al. (1984) *EMBO J.* 3(12): 2723-2730.  
30 207. Vernade et al. (1988) *J. Bacteriol.* 170: 5822-5829  
208. Wader et al., in *TOMATO TECHNOLOGY* 189-198 (Alan R. Liss, Inc. 1987)  
209. Wan & Lemaux (1994) *Plant Physiol.*, 104:3748  
210. Ward et al. (1993) *Plant Mol Biol* 22:361-366  
211. Waterhouse PM et al. (1998) *Proc Natl Acad Sci USA* 95:13959-64  
35 212. Watson et al. (1975) *J. Bacteriol* 123, 255-264  
213. Watson et al. (1985) *EMBO J* 4(2):277- 284  
214. Weeks et al. (1993) *Plant Physiol* 102:1077-1084  
215. WO 00/15815  
216. WO 00/26388  
40 217. WO 02/00900  
218. WO 84/02913  
219. WO 91/02071  
220. WO 91/13980  
221. WO 91/13991  
45 222. WO 92/06205  
223. WO 93/01294  
224. WO 93/18168  
225. WO 93/21334  
226. WO 94/00583

227. WO 94/00977  
228. WO 94/01999  
229. WO 94/48814  
230. WO 95/06722  
5 231. WO 95/15389  
232. WO 95/19443  
233. WO 95/23230  
234. WO 98/01576  
235. WO 98/45456  
10 236. WO 98/45461  
237. WO 99/16890  
238. WO 00/44895;  
239. WO 00/44914;  
240. WO 00/49035;  
15 241. WO 00/63364  
242. WO 00/68374;  
243. WO 97/41228;  
244. WO 98/45456  
245. WO 99/32619  
20 246. WO 99/53050;  
247. Xu et al. (1990) Chin J Bot 2(2):81-87  
248. Xu ZH et al. (1981) J. Exp. Bot. 32, 767-778.  
249. Zhou et al. (1992) Plant Cell Tissue Organ Culture, 30:78-83  
250. Zupan et al. (2000) Plant J 23(1):11-28  
25

## Claims

## WHAT IS CLAIMED IS:

- 5 1. A method for producing a transgenic Gramineae plant comprising the steps of:
- 10 (a) isolating a zygote from a Gramineae plant to be transformed in a way that said isolated zygote becomes substantially free from its naturally surrounding tissue,
- 15 (b) introducing a DNA composition comprising a genetic component into the genome of said Gramineae plant, wherein said introduction is mediated by Agrobacterium transformation into said isolated zygote;
- 20 (c) regenerating Gramineae plants from said zygotes which have received said genetic component; and
- (d) identifying a fertile, transgenic Gramineae plant whose genome has been altered through the stable introduction of said genetic component.
- 25 2. The method of claim 1, wherein the Gramineae plant is selected from the group consisting of wheat, maize and barley.
3. The method of claim 1 or 2, where in the Gramineae plant is a Triticum species.
- 30 4. The method of any of claim 1 to 3, wherein the Gramineae plant is regenerated from said isolated zygote by a method comprising co-cultivation of said isolated zygote and/or the zygotic embryo derived therefrom with a feeder system.
- 35 5. The method of any of claim 1 to 4, wherein the Gramineae plant is regenerated from said isolated zygote by a method comprising co-cultivation of said isolated zygote and/or the zygotic embryo derived therefrom with a culture of isolated immature pollen or pistils.
- 40 6. The method of any of claim 1 to 5, wherein the Gramineae plant is regenerated from said isolated zygote by a method comprising co-cultivation of said isolated zygote and/or the zygotic embryo derived therefrom with
- a) a culture of androgenetically developing barley pollen or
- b) a culture of wheat or barley pistils or
- c) any combination of a) and b).
- 45 7. The method of any of claim 1 to 6, wherein the zygotes and the feeder system are physically separated in a way to prevent mixing of the different cell types but to allow exchange of growth factors, proteins, media components, and other low molecular weight compounds.
8. The method of any of claim 1 to 7, wherein co-cultivation of the zygotes and the feeder system are employed already during Agrobacterium co-cultivation in a way

that the co-cultivation culture of the zygotes and Agrobacterium is physically separated from the feeder system in a way to prevent contact of the Agrobacteria with the feeder system but to allow exchange of growth factors, proteins, media components, and other low molecular weight compounds.

5

9. The method of any of claim 1 to 8, wherein said genetic component is transmitted through a complete sexual cycle of said transgenic Gramineae plant to its progeny, wherein said progeny does not comprise a selectable or screenable marker gene.

10

10. The method of any of claim 1 to 9, wherein said method does not comprise a step which leads to dedifferentiation of the zygote or the zygote-derived embryo.

15

11. The method of any of claim 1 to 10, wherein said genetic component comprises a expression cassette comprising a nucleic acid sequence operably linked to a promoter active in said Gramineae plant, wherein expression of said nucleic acid sequence confers a phenotypically distinguishable trait to said Gramineae plant.

20

12. The method of any of claim 1 to 11, wherein the pH of the medium used during co-cultivation of the isolated zygote with Agrobacterium is kept in a range from about 5.8 to about 6.0.

FIGURES

5

10

15

20

25

30




Fig. 1

**THIS PAGE BLANK (USPTO)**






Fig. 2

**THIS PAGE BLANK (USPTO)**




Fig. 3

**THIS PAGE BLANK (USPTO)**




Fig. 4

**THIS PAGE BLANK (USPTO)**

5

A

B

10

15

20

25

30




Fig. 5

**THIS PAGE BLANK (USPTO)**



## SEQUENCE LISTING

&lt;110&gt; BASF Plant Science GmbH

5 <120> METHODS FOR THE PRODUCTION OF STABLY TRANSFORMED, FERTILE  
 GRAMINEAE EMPLOYING AGROBACTERIUM-MEDIATED TRANSFORMATION OF ISOLATED  
 GRAMINEAE ZYGOTES

&lt;130&gt; PF 54757 - AE20030363

10

&lt;160&gt; 3

&lt;170&gt; PatentIn version 3.1

15

&lt;210&gt; 1

&lt;211&gt; 12739

&lt;212&gt; DNA

&lt;213&gt; synthetic: expression vector - binary vector pUGAB7

20

&lt;400&gt; 1

	tctagaacat ggtggagcac gacactctcg tctactccaa gaatatcaaa gatacagtct	60
	cagaagacca gagggctatt gagacttttc aacaaagggg aatatcgga aacctcctcg	120
	gattccattg cccagctatc tgtcacttca tcgaaaggac agtagaaaag gaagatggct	180
	tctacaaatg ccatcattgc gataaaggaa aggctatcgt tcaagatgcc tctaccgaca	240
25	gtggtcccaa agatggaccc ccaccacga ggaacatcgt ggaaaaagaa gacgttccaa	300
	ccacgtcttc aaagcaagtg gattgatgtg atatctccac tgacgtaagg gatgacgcac	360
	aatcccacta tccttcgcaa gacccttcct ctatataagg aagttcatit catttgga	420
	ggacctcgag tggccaccat gggcccagaa cgacgcccgg ccgacatccg ccgtgccacc	480
	gaggcggaca tgccggcggg ctgcaccatc gtcaaccact acatcgagac aagcacggtc	540
30	aacttcgcta ccgagccgca ggaaccgcag gaggggacgg acgacctcgt ccgtctgagg	600
	gagcgctatc cctggctcgt cgccgagggt gacggcgagg tcgcccggcat cgcctacgag	660
	ggcccctgga aggcacgcaa cgcctacgac tggacggcgg agtcgaccgt gtacgtctcc	720
	ccccgccacc agcggacggg actgggctcc acgctctaca cccacctgct gaagtccctg	780
	gaggcacagg gcttcaagag cgtggtcgtc gtcacggggc tgcccaacga cccgagcgtg	840
35	cgcatgacg aggcgctcgg atatgcccc cgcgcatgc tcgggggcgg cggttcaag	900
	cacgggaact ggcattgacgt gggtttctgg cagctggact tcagcctgcc ggtaccgccc	960
	cgtccgggtc tgcccgtcac cgagatctga gatcacgct tctagtccgc aaaaatcacc	1020
	agtctctctc taaaaatcta tctctctcta ttttctcca gaataatgtg tgagtagttc	1080
	ccagataagg gaattagggt tcttataggg tttcgctcat gtgttgagca tataagaaac	1140
40	ccttagtatg ttttgtatt tgtaaaatac ttctatcaat aaaatttcta attcctaaaa	1200
	ccaaaatcca gtgacctgca ggcattgcaag ctgatccact agaggccatg gcggccgctg	1260
	cgagcgatct agtaacatag atgacaccgc gcgcgataat ttatcctagt ttgcgcgcta	1320
	tattttgttt tctatcgctg attaaatgta taattgaggg actctaataca taaaaacca	1380
	tctcataaat aacgtcatgc attacatgtt aattattaca tgcttaacgt aattcaacag	1440
45	aaatttatg ataatcatcg caagaccggc aacaggattc aatcttaaga aactttattg	1500
	ccaaatggtt gaacgatcgg ggaaattcga gctcggtagc aattcccagag gctgtagccg	1560
	acgatgggtc gccaggagag ttgttgattc attgtttgcc tccctgctgc ggtttttcac	1620
	cgaagtcat gccagtcag cgtttttgca gcagaaaagc cgccgacttc ggtttgagggt	1680
	cgcgagtga gatccctttc ttgttaccgc caacgcgcaa tatgccttgc gaggtcgcaa	1740

	aatcggcgaa	attccatacc	tggtcaccga	cgacggcgct	gacgcgatca	aagacgcggt	1800
	gatacatatc	cagccatgca	cactgatact	cttcactcca	catgtcgggtg	tacattgagt	1860
	gcagcccggc	taacgtatcc	acgccgtatt	cggtgatgat	aatcggctga	tgcagtttct	1920
	cctgccaggc	cagaagttct	ttttccagta	ccttctctgc	cgtttccaaa	tcgccgcttt	1980
5	ggacatacca	tccgtaataa	cggttcaggc	acagcacatc	aaagagatcg	ctgatgggtat	2040
	cggtgtgagc	gtcgcagAAC	attacattga	cgaggtgat	cggacgcgtc	gggtcgagtt	2100
	tacgcgttgc	ttccgccagt	ggcgcgaaat	attcccgtgc	accttgcgga	cggttatccg	2160
	gttcggttggc	aatactccac	atcaccacgc	ttgggtgggt	tttgtcacgc	gctatcagct	2220
	ctttaatcgc	ctgtaagtgc	gcttgctgag	tttcccgtt	gactgcctct	tcgctgtaca	2280
10	gttcttttcgg	cttggttgccc	gcttcgaaac	caatgcctaa	agagagggtta	aagccgacag	2340
	cagcagtttc	atcaatcacc	acgatgccat	gttcatctgc	ccagtcgagc	atctcttcag	2400
	cgtaagggtta	atgCGaggta	cggtaggagt	tggccccaat	ccagtccatt	aatgcgtgggt	2460
	cgtgcaccat	cagcacgtta	tcgaatcctt	tgccacgcaa	gtccgcacat	tcatgacgac	2520
	caaagccagt	aaagtagaac	ggtttgtgggt	taatcaggaa	ctggttgccc	ttcactgcca	2580
15	ctgaccggat	gccgacgcga	agcgggtaga	tatcacactc	tgtctggctt	ttggctgtga	2640
	cgcacagttc	atagagataa	ccttcacccg	gttgccagag	gtgcggattc	accacttgca	2700
	aagtcccgt	agtgccttgt	ccagttgcaa	ccacctgttg	atccgcatca	cgcagttcaa	2760
	cgctgacatc	accattggcc	accacctgcc	agtcaacaga	cgcgtgggtta	cagtcttgcg	2820
	cgacatgcgt	caccacgggtg	atatactcca	cccagggtgtt	cggcgtgggtg	tagagcatta	2880
20	cgctgcgatg	gattccggca	tagttaaaga	aatcatggaa	gtaagactgc	tttttcttgc	2940
	cgttttctgc	ggtaatcacc	attcccggcg	ggatagtctg	ccagttcagt	tcgttgttca	3000
	cacaaacgggt	gataccagct	gcacatcaac	aaattttgggt	catatattag	aaaagttata	3060
	aattaaaata	tacacactta	taaactacag	aaaagcaatt	gctatatact	acattctttt	3120
	attttgaaaa	aaatatattga	aatattatat	tactactaat	taatgataat	tattatatat	3180
25	atatcaaagg	tagaagcaga	aacttacgta	cacttttccc	ggcaataaca	tacggcgtga	3240
	catcggcttc	aaatggcgta	tagccgccct	gatgctccat	cacttcctga	ttattgaccc	3300
	acactttgcc	gtaatgagtg	accgcacgca	aacgcagcac	gatacgctgg	cctgcccAAC	3360
	ctttcgggtat	aaagacttcg	cgctgatacc	agacgttgcc	cgcataatta	cgaatatctg	3420
	catcggcgaa	ctgatcgtta	aaactgcctg	gcacagcaat	tgcccggctt	tcttgtaacg	3480
30	cgctttccca	ccaacgctga	ccaattccac	agttttcgcg	atccagactg	aatgcccaca	3540
	ggccgctcag	ttttttgatt	tcacgggttg	gggtttctac	aggacgtaac	ataagggact	3600
	gacgtcgaat	ccactagtag	ctgcagaagt	aacaccaaac	aacagggtga	gcatcgacaa	3660
	aagaaacagt	accaagcaaa	taaatagcgt	atgaaggcag	ggctaaaaaa	atccacatat	3720
	agctgctgca	tatgccatca	tccaagtata	tcaagatcaa	aataattata	aaacatactt	3780
35	gtttattata	atagatagggt	actcaagggt	agagcatatg	aatagatgct	gcatatgcca	3840
	tcatgtatat	gcatcagtaa	aaccacacatc	aacatgtata	cctatcctag	atcgatcccg	3900
	tctgcggaac	ggctagagcc	atcccaggat	tcccaaaga	gaaacactgg	caagttagca	3960
	atcagaacgt	gtctgacgta	caggtcgcac	ccgtgtacga	acgctagcag	cacggatcta	4020
	acacaaacac	ggatctaaca	caaacatgaa	cagaagtaga	actaccgggc	cctaaccatg	4080
40	gaccggaacg	ccgatctaga	gaaggtagag	aggggggggg	ggggaggacg	agcggcgtag	4140
	cttgaagcgg	aggtgccgac	gggtggattt	gggggagatc	tgggtgtgtg	tgtgtgcgct	4200
	ccgaacaaca	cgaggttggg	gaaagaggggt	gtggaggggg	tgtctattta	ttacggcggg	4260
	cgaggaaggg	aaagcgaagg	agcggtgagg	aaggaatccc	ccgtagctgc	cggtgccgtg	4320
	agaggaggag	gaggccgcct	gccgtgccgg	ctcacgtctg	ccgctccgcc	acgcaatttc	4380
45	tggatgccga	cagcggagca	agtccaacgg	tggagcggaa	ctctcgagag	gggtccagag	4440
	gcagcgacag	agatgccgtg	ccgtctgctt	cgcttgcccc	gacgcgacgc	tgtgtggttcg	4500
	ctggttggtg	tccgttagac	tcgtcgatcg	acggcgcttta	acaggctggc	attatctact	4560
	cgaacaaga	aaaatgtttc	cttagttttt	ttaatttctt	aaagggtatt	tgtttaattt	4620
	ttagtcactt	tattttattc	tattttatat	ctaaattatt	aaataaaaaa	actaaaatag	4680

	agtttttagtt	ttcttaatttt	agaggctaaa	atagaataaaa	atagatgtac	taaaaaaatt	4740
	agtctataaaa	aaccattaac	cctaaaccct	aatggatgt	actaataaaa	tggatgaagt	4800
	attatatagg	tgaagctatt	tgcaaaaaaa	aaggagaaca	catgcacact	aaaaagataa	4860
	aactgtagag	tcctgttgct	aaaataactca	attgtccttt	agaccatgtc	taactgttca	4920
5	tttatatgat	tctctaaaac	actgatatta	ttgtagtact	atagattata	ttattcgtag	4980
	agtaaagttt	aaatatatgt	ataaagatag	ataaaactgca	cttcaaaca	gtgtgacaaa	5040
	aaaaatatgt	ggtaattttt	tataacttag	acatgcaatg	ctcattatct	ctagagaggg	5100
	gcacgaccgg	gtcacgctgc	actgcagcct	agtaaggcct	taagggccag	atcttgggcc	5160
	cggtagccga	tcagattgtc	gtttcccgcc	ttcggtttaa	actatcagt	tttgacagga	5220
10	tatattggcg	ggtaaaccta	agagaaaaga	gcgtttatta	gaataatcgg	atatttaaaa	5280
	gggcgtgaaa	aggtttatcc	gttcgtccat	ttgtatgtgc	atgccaacca	cagggttccc	5340
	ctcgggagtg	cttggcattc	cgtgcgataa	tgacttctgt	tcaaccaccc	aaacgtcgga	5400
	aagcctgacg	acggagcagc	attccaaaaa	gatcccttgg	ctcgtctggg	tcggctagaa	5460
	ggtcgagtg	gctgctgtgg	cttgatccct	caacgcggtc	gcggacgtag	cgcagcgccg	5520
15	aaaaatcctc	gatcgcaaat	ccgacgctgt	cgaaaagcgt	gatctgcttg	tcgctctttc	5580
	ggccgacgtc	ctggccagtc	atcacgcgcc	aaagttccgt	cacaggatga	tctggcgcca	5640
	gttgctggat	ctcgccttca	atccgggtct	gtggcgggaa	ctccacgaaa	atatccgaac	5700
	gcagcaagat	atcgcggtgc	atctcggtct	tgccctgggca	gtcgccgccg	acgccgttga	5760
	tgtggacgcc	gaaaaggatc	taggtgaaga	tcctttttga	taatctcatg	acaaaaatcc	5820
20	cttaacgtga	gttttcgttc	caactgagcgt	cagaccccg	agaaaagatc	aaaggatctt	5880
	cttgagatcc	tttttttctg	cgcgtaatct	gctgcttgca	aacaaaaaaa	ccaccgctac	5940
	cagcgggtgg	ttgtttgccc	gatcaagagc	taccaactct	ttttccgaag	gtaactggct	6000
	tcagcagagc	gcagatacca	aatactgttc	ttctagtgt	gccgtagtta	ggccaccact	6060
	tcaagaactc	tgtagcaccg	cctacatacc	tcgctctgct	aatcctgtta	ccagtggctg	6120
25	ctgccagtg	cgataagtcg	tgtcttaccg	ggttggactc	aagacgatag	ttaccggata	6180
	aggcgcagcg	gtcgggctga	acgggggggt	cgtgcacaca	gcccagcttg	gagcgaacga	6240
	cctacaccga	actgagatac	ctacagcgtg	agctatgaga	aagcgccacg	cttcccgaag	6300
	ggagaaaggc	ggacaggtat	ccggtaaagc	gcagggctcg	aacaggagag	cgcacgaggg	6360
	agcttccagg	gggaaacgcc	tggtatcttt	atagtcctgt	cgggtttcgc	cacctctgac	6420
30	ttgagcgtcg	atttttgtga	tgctcgtcag	gggggcggag	cctatggaaa	aacgccagca	6480
	acgcggcctt	tttacgggtc	ctggcctttt	gctggccttt	tgctcacatg	ttctttcctg	6540
	cgttatcccc	tgattctgtg	gataaccgat	taccgccttt	gagtgagctg	ataccgctcg	6600
	ccgcagccga	acgaccgagc	gcagcagctc	agtgcgagag	gaagcggaag	agcgcctgat	6660
	gcggatattt	ctccttacgc	atctgtgcgg	tatttcacac	cgcatatgg	gcactctcag	6720
35	tacaatctgc	tctgatgccg	catagttaag	ccagtataca	ctccgctatc	gctacgtgac	6780
	tgggtcatgg	ctgcgccccg	acacccgcca	acacccgctg	acgcgcccctg	acgggcttgt	6840
	ctgctcccgg	catccgctta	cagacaagct	gtgaccgtct	ccgggagctg	catgtgtcag	6900
	aggttttcac	cgtcatcacc	gaaacgcgcg	aggcaggggt	acgtcgaggt	cgatccaacc	6960
	cctccgctgc	tatagtgcag	tcggcttctg	acgttcagtg	cagccgtctt	ctgaaaacga	7020
40	catgtcgcac	aagtcctaag	ttacgcgaca	ggctgccgcc	ctgccctttt	cctggcgctt	7080
	tcttgtcgcg	tgtttttagtc	gcataaagta	gaataactgc	gactagaacc	ggagacatta	7140
	cgccatgaac	aagagcgccg	ccgctggcct	gctgggctat	gcccgcgtca	gcaccgacga	7200
	ccaggacttg	accaaccaac	gggcccgaact	gcacgcggcc	ggctgcacca	agctgttttc	7260
	cgagaagatc	accggcacca	ggcgcgaccg	cccggagctg	gccaggatgc	ttgaccacct	7320
45	acgccctggc	gacgttgtga	cagtgaccag	gctagaccgc	ctggcccgcga	gcacccgcga	7380
	cctactggac	attgccgagc	gcataccagga	ggccggcgcg	ggcctgcgta	gcctggcaga	7440
	gccgtggggc	gacaccacca	cgccggccgg	ccgcatgggt	ttgaccgtgt	tcgccggcat	7500
	tgccgagttc	gagcgttccc	taatcatcga	ccgcaccccg	agcggggcgcg	aggccgccaa	7560
	ggcgcgaggg	gtgaagtttg	gccccgcgcc	taccctcacc	ccggcacaga	tcgcgcacgc	7620

	ccgcgagctg	atcgaccagg	aaggccgcac	cgtgaaagag	gcggctgcac	tgcttggcgt	7680
	gcacgcgtcg	accctgtacc	gcgcacttga	gcgcagcgag	gaagtgcgc	ccaccgaggc	7740
	caggcgggcg	ggtgccttcc	gtgaggacgc	attgaccgag	gccgacgccc	tgggcgggcg	7800
	cgagaatgaa	cgccaagagg	aacaagcatg	aaaccgcacc	aggacggcca	ggacgaaccg	7860
5	tttttcatta	ccgaagagat	cgaggcggag	atgatcgcg	ccgggtacgt	gttcgagccg	7920
	cccgcgcacg	tctcaaccgt	gcggctgcat	gaaatcctgg	ccggtttgtc	tgatgccaag	7980
	ctcgcgccct	ggccggcgag	cttgcccgct	gaagaaaccg	agcgccgccc	tctaaaaagg	8040
	tgatgtgtat	ttgagtaaaa	cagcttgctg	catgcggtcg	ctgcgtatat	gatgcgatga	8100
	gtaaataaac	aaatacgcaa	ggggaacgca	tgaaggttat	cgctgtactt	aaccagaaaag	8160
10	gcgggtcagg	caagacgacc	atcgcaaccc	atctagcccc	cgccctgcaa	ctcgccgggg	8220
	ccgatgttct	gttagtcgat	tccgatcccc	agggcagtg	ccgcgattgg	gcggccgtgc	8280
	gggaagatca	accgctaacc	gttgtcggca	tcgaccgccc	gacgattgac	cgcgacgtga	8340
	aggccatcgg	ccggcgcgac	ttcgtagtga	tcgacggagc	gccccaggcg	gcggacttgg	8400
	ctgtgtccgc	gatcaaggca	gccgacttcg	tgctgattcc	ggtgcagcca	agcccttacg	8460
15	acatatgggc	caccgcccac	ctggtggagc	tggttaagca	gcgcattgag	gtcacggatg	8520
	gaaggctaca	agcggccttt	gtcgtgtcgc	gggcgatcaa	aggcacgcgc	atcggcggtg	8580
	aggttgccga	ggcgctggcc	gggtacgagc	tgcccattct	tgagtcccgt	atcacgcagc	8640
	gcgtgagcta	cccaggcact	gccgcgcgcg	gcacaaccgt	tcttgaatca	gaacccgagg	8700
	gcgacgctgc	ccgcgaggtc	caggcgctgg	ccgctgaaat	taaatcaaaa	ctcatttgag	8760
20	ttaatgaggt	aaagagaaaa	tgagcaaaaag	cacaaacacg	ctaagtgccg	gccgtccgag	8820
	cgcacgcagc	agcaaggctg	caacgttggc	cagcctggca	gacacgccag	ccatgaagcg	8880
	ggtcaacttt	cagttgcccg	cggaggatca	caccaagctg	aagatgtacg	cggtagccca	8940
	aggcaagacc	attaccgagc	tgctatctga	atacatcgcg	cagctaccag	agtaaataag	9000
	caaatagaata	aatgagtaga	tgaatttttag	cggctaaagg	aggcggcag	gaaaatacaag	9060
25	aacaaccagg	caccgacgcc	gtggaatgcc	ccatgtgtgg	aggaacgggc	ggttggccag	9120
	gcgtaagcgg	ctgggttgtc	tgccggccct	gcaatggcac	tggaaccccc	aagcccgagg	9180
	aatcggcggtg	agcggtcgca	aaccatccgg	cccggtagaa	atcggcgcg	cgctgggtga	9240
	tgacctgggtg	gagaagttga	aggcgccgca	ggccgcccag	cggcaacgca	tcgaggcaga	9300
	agcacgcccc	ggtgaatcgt	ggcaagcggc	cgtgatcgca	atccgcaaag	aatcccggca	9360
30	accgcccggca	gccggtgcgc	cgtcgattag	gaagccgccc	aaggcgacg	agcaaccaga	9420
	ttttttcggt	ccgatgctct	atgacgtggg	caccgcgcg	agtcgcagca	tcattggacgt	9480
	ggccgttttc	cgtctgtcga	agcgtgaccg	acgagctggc	gaggtgatcc	gctacgagct	9540
	tccagacggg	cacgtagagg	tttccgcagg	gccggccggc	atggcgagt	tgtgggatta	9600
	cgacctggta	ctgatggcgg	tttcccatct	aaccgaatcc	atgaaccgat	accgggaagg	9660
35	gaagggagac	aagcccggcc	gcgtgttccg	tccacacgtt	gcggacgtac	tcaagttctg	9720
	ccggcgagcc	gatggcggaa	agcagaaaag	cgacctggta	gaaacctgca	ttcggttaaa	9780
	caccacgcac	gttgccatgc	agcgtacgaa	gaaggccaag	aacggccgcc	tggtgacggt	9840
	atccgagggg	gaagccttga	ttagccgcta	caagatcgta	aagagcgaaa	ccgggcccgc	9900
	ggagtacatc	gagatcgagc	tagctgattg	gatgtaccgc	gagatcacag	aaggcaagaa	9960
40	cccgacgtg	ctgacggttc	accccgatta	ctttttgatc	gatcccggca	tcggccgttt	10020
	tctctaccgc	ctggcacgcc	gcgccgcagg	caaggcagaa	gccagatggt	tgttcaagac	10080
	gatctacgaa	cgcagtggca	gcgccggaga	gttcaagaag	ttctgtttca	ccgtgcgcaa	10140
	gctgatcggg	tcaaatgacc	tgccggagta	cgatttgaag	gaggaggcgg	ggcaggctgg	10200
	cccgatccta	gtcatgcgct	accgcaacct	gatcgagggc	gaagcatccg	ccggttccta	10260
45	atgtacggag	cagatgctag	ggcaaatgac	cctagcaggg	gaaaaagggtc	gaaaaaggctc	10320
	ctttcctgtg	gatagcacgt	acattgggaa	cccaaagccg	tacattggga	accggaaccc	10380
	gtacattggg	aacccaaagc	cgtacattgg	gaaccggtca	cacatgtaag	tgactgatata	10440
	aaaagagaaa	aaaggcgatt	tttccgccta	aaactcttta	aaacttatta	aaactcttaa	10500
	aaccgcgctg	gcctgtgcat	aactgtctgg	ccagcgcaca	gccgaagagc	tgcaaaaagc	10560

gcctaccctt cggtcgctgc gctccctacg ccccgccgct tcgcgctcggc ctatcgcggc 10620  
 cgctggccgc tcaaaaatgg ctggcctacg gccaggcaat ctaccagggc gcggaacaagc 10680  
 cgcgcgctcg cactcgacc gccggcgccc acatcaaggc accggtgggt atgcctgacg 10740  
 atgcgtggag accgaaacct tgcgctcggt cgccagccag gacagaaatg cctcgacttc 10800  
 5 gctgctgccc aagggtgccc ggtgacgcac accgtggaaa cggatgaagg cacgaacca 10860  
 gtggacataa gcctgttcgg ttcgtaagct gtaatgcaag tagcgtatgc gctcacgcaa 10920  
 ctggtccaga accttgaccg aacgcagcgg tggtaacggc gcagtggcgg ttttcatggc 10980  
 ttgttatgac tgtttttttg gggtagctc tatgcctcgg gcatccaagc agcaagcgcg 11040  
 ttacgccgtg ggtcgatggt tgatgttatg gagcagcaac gatgttacgc agcagggcag 11100  
 10 tcgccctaaa acaaagttaa acatcatgag ggaagcgggt atcgccgaag tatcgactca 11160  
 actatcagag gtagttggcg tcatcgagcg ccattctcga ccgacgttgc tggccgtaca 11220  
 tttgtacggc tccgcagtgg atggcgccct gaagccacac agtgatattg atttgctggg 11280  
 tacggtgacc gtaaggcttg atgaaacaac gcggcgagct ttgatcaacg accttttgga 11340  
 aacttcgggt tcccctggag agagcgagat tctccgcgct gtagaagtca ccattgttgt 11400  
 15 gcacgacgac atcattccgt ggcgttatcc agctaagcgc gaactgcaat ttggagaatg 11460  
 gcagcgcaat gacattcttg caggtatctt cgagccagcc acgatcgaca ttgatctggc 11520  
 tatcttgctg acaaaagcaa gagaacatag cgttgccctg gtaggtccag cggcggagga 11580  
 actctttgat cgggttcctg aacaggatct atttgaggcg ctaaataaaa ccttaacgct 11640  
 atggaactcg ccgcccact gggctggcga tgagcgaaat gtagtgctta cgttgtcccg 11700  
 20 catttggtac agcgcagtaa ccggcaaat cgcgccgaag gatgtcgctg ccgactgggc 11760  
 aatggagcgc ctgccggccc agtatcagcc cgtcatactt gaagctagac aggcttatct 11820  
 tggacaagaa gaagatcgct tggcctcgcg cgcagatcag ttggaagaat ttgtccacta 11880  
 cgtgaaaggc gagatcacca aggtagtcgg caaataatgt ctaacaattc gttcaagccg 11940  
 acgccgcttc gcggcgcggc ttaactcaag cgttagatgc actaagcaca taattgctca 12000  
 25 cagccaaact atcaggtcaa gtctgctttt attattttta agcgtgcata ataagcccta 12060  
 cacaaattgg gagatatatc atgaaaggct ggctttttct tgttatcgca atagttggcg 12120  
 aagtaatcgc aacatagctt gcttggtcgt tccgcgtgaa cgtcggctcg attgtacctg 12180  
 cgttcaaata ctttgcgatc gtgttgcgcg cctgcccggt gcgtcggctg atctcacgga 12240  
 tcgactgctt ctctcgcaac gccatccgac ggatgatggt taaaagtccc atgtggatca 12300  
 30 ctccgttgcc ccgtcgctca ccgtgttggg gggaagggtc acatggctca gttctcaatg 12360  
 gaaattatct gcctaaccgg ctgagttctg cgtagaaacc aacatgcaag ctccaccggg 12420  
 tgcaaaagcg cagcggcggc aggatatatt caattgtaaa tggcttcatg tccgggaaat 12480  
 ctacatggat cagcaatgag tatgatggtc aatatggaga aaaagaaaga gtaattacca 12540  
 attttttttc aattcaaaaa tgtagatgtc cgcagcgtta ttataaaatg aaagtacatt 12600  
 35 ttgataaaac gacaaattac gatccgctgt atttataggc gaaagcaata aacaaattat 12660  
 tctaattcgg aaatctttat ttcgacgtgt ctacattcac gtccaaatgg gggcttagat 12720  
 gagaaacttc acgatcggc 12739

<210> 2

40

<211> 11195

<212> DNA

<213> synthetic: expression vector - binary vector pSUNUbi-GFP

45

<400> 2

aatattcaaa caaacacata cagcgcgact tatcatggac atacaaatgg acgaacggat 60  
 aaaccttttc acgccctttt aaatatccga ttattctaata aaacgctctt ttctcttagg 120  
 tttaccgcgc aatatatcct gtcaaacact gatagtttaa actgaaggcg ggaaacgaca 180  
 atcagatcta gtaggaaaca gctatgacca tgattacgcc aagctggccg cgtcgagcga 240

	tctagtaaca	tagatgacac	cgcgcgcgat	aattttatcct	agtttgcgcg	ctatatatttg	300
	ttttctatcg	cgtattaaat	gtataattgc	gggactctaa	tcataaaaac	ccatctcata	360
	aataacgtca	tgcattacat	gttaattatt	acatgcttaa	cgtaattcaa	cagaaattat	420
	atgataatca	tgcgaagacc	ggcaacagga	ttcaatctta	agaaacttta	ttgccaaatg	480
5	tttgaacgat	cggggaaatt	cgagtcgact	ctagaggatc	cccggccgcc	ccgactctag	540
	tgcgattact	tgtacagctc	gtccatgcog	agagtgatcc	cggcgccggt	cacgaactcc	600
	agcaggacca	tgtgatcgcg	cttctcggtg	gggtctttgc	tcagggcgga	ctgggtgctc	660
	aggtagtggt	tgtcgggcag	cagcacgggg	ccgtcgccga	tgggggtggt	ctgctggtag	720
	tggtcggcga	gctgcacgct	gccgtcctcg	atgttggtggc	gggtcttgaa	gttcaccttg	780
10	atgccgttct	tctgcttgct	ggcggtgata	tagaccttgt	ggctggtgta	gttgactcc	840
	agcttgctgc	ccaggatggt	gccgtcctcc	ttgaagtcga	tgcccttcag	ctcgatgcgg	900
	ttcaccaggg	tgtcgccctc	gaacttcacc	tcggcgccgg	tcttgtagtt	gccgtcgtcc	960
	ttgaagaaga	tgggtgcgctc	ctggacgtag	ccttcgggca	tggcggactt	gaagaagtcg	1020
	tgctgcttca	tgtggtcggg	gtagcggggc	aagcactgca	cgcgtaggt	caagggtggtc	1080
15	acgagggtgg	gccagggcac	gggcagcttg	ccggtggtgc	agatgaactt	cagggtcagc	1140
	ttgccgtagg	tggcatcgcc	ctcgccctcg	ccggacacgc	tgaacttggt	gccgtttacg	1200
	tcgccgtcca	gctcgaccag	gatgggcacc	accccggtga	acagctcttc	gcccttgctc	1260
	accatggtgg	ctttaccaac	agtaccgga	tgccaagctt	acttagatcg	cagatctact	1320
	agtatgatgg	gtaccggggg	gatccgatat	cgcccgggct	gcagaagtaa	caccaacaa	1380
20	cagggtgagc	atcgacaaaa	gaaacagtac	caagcaaata	aatagcgat	gaaggcaggg	1440
	ctaaaaaat	ccacatatag	ctgctgcata	tgccatcatc	caagtatatc	aagatcaaaa	1500
	taattataaa	acatacttgt	ttattataat	agataggtag	tcaagggttag	agcatatgaa	1560
	tagatgctgc	atatgccatc	atgtatatgc	atcagtaaaa	cccacatcaa	catgtatacc	1620
	tatcctagat	cgatatattcc	atccatctta	aactcgtaac	tatgaagatg	tatgacacac	1680
25	acatacagtt	ccaaaattaa	taaatacacc	aggtagtttg	aaacagtatt	ctactccgat	1740
	ctagaacgaa	tgaacgaccg	cccaaccaca	ccacatcatc	acaaccaagc	gaacaaaaag	1800
	catctctgta	tatgcatcag	taaaaccgcg	atcaacatgt	atacctatcc	tagatcgata	1860
	tttccatcca	tcatcttcaa	ttcgtaacta	tgaatatgta	tggcacacac	atacagatcc	1920
	aaaattaata	aatccaccag	gtagtttgaa	acagaattct	actccgatct	agaacgaccg	1980
30	cccaaccaga	ccacatcatc	acaaccaaga	caaaaaaaag	catgaaaaga	tgaccgcaca	2040
	aacaagtgca	cggcatatat	tgaataaaag	gaaaaggggca	aaccaaaacc	tatgcaacga	2100
	aacaaaaaaa	atcatgaaat	cgatcccgtc	tgcggaacgg	ctagagccat	cccaggattc	2160
	cccaaagaga	aacactggca	agtttagcaat	cagaacgtgt	ctgacgtaca	ggtcgcatcc	2220
	gtgtacgaac	gctagcagca	cggatctaac	acaaacacgg	atctaacaca	aacatgaaca	2280
35	gaagtagaac	taccggggcc	taaccatgga	ccggaacgcc	gatctagaga	aggtagagag	2340
	gggggggggg	ggaggacgag	cggcgctacct	tgaagcggag	gtgccgacgg	gtggatttg	2400
	gggagatctg	gttggtgtgt	tgtgcgctcc	gaacaacacg	aggttgggga	aagagggtgt	2460
	ggagggggtg	tctattttatt	acggcggggc	aggaagggaa	agcgaaggag	cgggtgggaa	2520
	ggaatcccc	gtagctgccg	tgccgtgaga	ggaggaggag	gccgcctgcc	gtgccggctc	2580
40	acgtctgcg	ctccgccacg	caattttctg	atgccgacag	cggagcaagt	ccaacgggtg	2640
	agcggaaactc	tgcgaggggg	tccagaggca	gcgacagaga	tgccgtgccg	tctgcttcgc	2700
	ttggcccgac	gcgacgctgc	tgggtcgctg	gttggtgtcc	gttagactcg	tcgacggcgt	2760
	ttaacaggct	ggcattatct	actcgaaaca	agaaaaatgt	ttccttagtt	tttttaattt	2820
	cttaaagggt	atttgtttaa	tttttagtca	ctttatttta	ttctatttta	tatctaaatt	2880
45	attaaataaa	aaaactaaaa	tagagtttta	gttttcttaa	tttagaggct	aaaatagaat	2940
	aaaatagatg	tactaaaaaa	attagtctat	aaaaaccatt	aaccctaaac	cctaaatgga	3000
	tgtactaata	aaatggatga	agtattatat	aggtagaagt	atttgcaaaa	aaaaaggaga	3060
	acacatgcac	actaaaaaga	taaaactgta	gagtcctgtt	gtcaaaaatac	tcaattgtcc	3120
	tttagaccat	gtctaactgt	tcattttatat	gattctctaa	aacactgata	ttattgtagt	3180

	actatagatt	atattattcg	tagagtaaag	tttaaata	tgtataaaga	tagataaact	3240
	gcacttcaaa	caagtgtgac	aaaaaaaaata	tgtggtaatt	ttttataact	tagacatgca	3300
	atgctcatta	tctctagaga	gggggcacgac	cgggtcacgc	tgcactgcag	gaattcgatg	3360
	ggctcgaggt	accgagctcg	gtacccactg	gatttttggt	ttaggaatta	gaaattttat	3420
5	tgatagaagt	attttcaaaa	tacaaatata	tactaagggt	ttcttatatg	ctcaacacat	3480
	gagcgaaacc	ctataagaac	cctaattccc	ttatctggga	actactcaca	cattattata	3540
	gagagagata	gatttgtaga	gagagactgg	tgatttcagc	ggcatgcctg	caggctcgact	3600
	cagatctggg	taactggcct	aactggcctt	ggaggagctg	gcaactcaaa	atccctttgc	3660
	caaaaaccaa	catcatgcca	tccaccatgc	ttgtatccag	ctgcgcgcaa	tgtaccccg	3720
10	gctgtgtatc	ccaaagcctc	atgcaacct	acagatggat	cgtttggaag	gcctataaca	3780
	gcaaccacag	acttaaaacc	ttgcgcctcc	atagacttaa	gcaaagtgtg	gtacaatgtg	3840
	gatcctaggc	ccaacctttg	atgcctatgt	gacacgtaaa	cagtactctc	aactgtccaa	3900
	tcgtaagcgt	tcctagcctt	ccagggccca	gcgtaagcaa	taccagccac	aacaccctca	3960
	acctcagcaa	ccaaccaagg	gtatctatct	tgcaacctct	ctagatcctc	aatccactct	4020
15	tgtggtgttt	gtggctctgt	cctaaagttc	actgtagacg	tctcaatgta	atgggttaacg	4080
	atatcacaaa	ccgcggccat	atcagctgct	gtagctggcc	taatctcaac	tggctcctc	4140
	tccggagaca	tgtcgactct	agaggatccc	ccgggctgca	gaagtaacac	caaacaacag	4200
	ggtgagcatc	gacaaaagaa	acagtaccaa	gcaataaat	agcgtatgaa	ggcagggtca	4260
	aaaaaatcca	catatagctg	ctgcatatgc	catcatccaa	gtatatcaag	atcaaaataa	4320
20	ttataaaaaca	tacttgttta	ttataataga	taggtactca	aggttagagc	atatgaatag	4380
	atgctgcata	tgccatcatg	tatatgcata	agtaaaaccc	acatcaacat	gtatacctat	4440
	cctagatcga	tatttccatc	catcttaaac	tcgtaactat	gaagatgtat	gacacacaca	4500
	tacagttcca	aaattaataa	atacaccagg	tagtttgaaa	cagtattcta	ctccgatcta	4560
	gaacgaatga	acgaccgccc	aaccacacca	catcatcaca	accaagcgaa	caaaaagcat	4620
25	ctctgtatat	gcatcagtaa	aaccgcgcatc	aacatgtata	cctatcctag	atcgatattt	4680
	ccatccatca	tcttcaattc	gtaactatga	atatgtatgg	cacacacata	cagatccaaa	4740
	attaataaat	ccaccaggta	gtttgaaaca	gaattctact	ccgatctaga	acgaccgccc	4800
	aaccagacca	catcatcaca	accaagacaa	aaaaaagcat	gaaaagatga	cccgcacaa	4860
	aagtgcacgg	catatattga	aataaaggaa	aagggcacaa	caaaccctat	gcaacgaaac	4920
30	aaaaaaaatc	atgaaatcga	tcccgctctg	ggaacggcta	gagccatccc	aggattcccc	4980
	aaagagaaac	actggcaagt	tagcaatcag	aacgtgtctg	acgtacaggt	cgcacccgtg	5040
	tacgaacgct	agcagcacgg	atctaacaca	aacacggatc	taacacaaac	atgaacagaa	5100
	gtagaactac	cgggccctaa	ccatggaccg	gaacgccgat	ctagagaagg	tagagagggg	5160
	ggggggggga	ggacgagcgg	cgtaccttga	agcggagggtg	ccgacgggtg	gatttggggg	5220
35	agatctgggt	gtgtgtgtgt	gcgctccgaa	caacacgagg	ttggggaaag	aggggtgtgga	5280
	gggggtgtct	atattattacg	gcgggcgagg	aagggaaagc	gaaggagcgg	tgggaaagga	5340
	atcccccgta	gctgccgtgc	cgtgagagga	ggaggaggcc	gcctgccgtg	ccggctcacg	5400
	tctgccgctc	cgccacgcaa	tttctggatg	ccgacagcgg	agcaagtcca	acgggtggagc	5460
	ggaactctcg	agaggggtcc	agaggcagcg	acagagatgc	cgtgccgtct	gcttcgcttg	5520
40	gcccgcagcg	acgctgctgg	ttcgctgggt	ggtgtccgtt	agactcgtcg	acggcggttta	5580
	acaggctggc	attatctact	cgaaacaaga	aaaatgtttc	cttagttttt	ttaattttctt	5640
	aaagggtatt	tgttttaattt	ttagtcactt	tattttattc	tattttatat	ctaaattatt	5700
	aaataaaaaa	actaaaatag	agtttttagtt	ttcttaattt	agaggctaaa	atagaataaa	5760
	atagatgtac	taaaaaaatt	agtctataaa	aaccattaac	cctaaaccct	aatggatgt	5820
45	actaataaaa	tggatgaagt	attatatagg	tgaagctatt	tgcaaaaaaa	aaggagaaca	5880
	catgcacact	aaaaagataa	aactgtagag	tcctgttgtc	aaaataactca	attgtccttt	5940
	agaccatgtc	taactgttca	tttatatgat	tctctaaaac	actgatattt	ttgtagtact	6000
	atagattata	ttattcgtag	agtaaagttt	aaatatatgt	ataaagatag	ataaactgca	6060
	cttcaaaca	gtgtgacaaa	aaaaatatgt	ggtaattttt	tataacttag	acatgcaatg	6120



	ctcattatct	ctagagaggg	gcacgaccgg	gtcacgctgc	actgcaggaa	ttcgatatca	6180
	agctggctga	gtggctcctt	caacgttgcg	gttctgtcag	ttccaaacgt	aaaacggctt	6240
	gtcccgcgtc	atcggcgggg	gtcataacgt	gactccctta	attctccgct	catgatcaga	6300
	ttgtcgtttc	ccgccttcag	tttaaaactat	cagtgtttga	caggatcact	gcttggtaat	6360
5	aattgtcatt	agattgtttt	tatgcataga	tgcactcgaa	atcagccaat	tttagacaag	6420
	tatcaaacgg	atgttaattc	agtacattaa	agacgtccgc	aatgtgttat	taagttgtct	6480
	aagcgtcaat	ttgttttacac	cacaatatat	cctgccacca	gccagccaac	agctccccga	6540
	ccggcagctc	ggcacaaaat	caccacgcgt	ctaaaaagg	gatgtgtatt	tgagtaaaac	6600
	agcttgcgctc	atgcggctgc	tgcgtatatg	atgcgatgag	taaataaaca	aatacgcaag	6660
10	gggaacgcat	gaagggttatc	gctgtactta	accagaaagg	cgggtcaggc	aagacgacca	6720
	tcgcaaccca	tctagcccg	gccctgcaac	tcgccggggc	cgatgttctg	ttagtcgatt	6780
	ccgatcccca	gggcagtgcc	cgcgattggg	cggccgtg	ggaagatcaa	ccgctaaccg	6840
	ttgtcggcat	cgaccgccc	acgattgacc	gcgacgtgaa	ggccatcggc	cggcgcgact	6900
	tcgtagtgat	cgacggagcg	ccccaggcgg	cggacttg	tgtgtccg	atcaaggcag	6960
15	ccgacttcgt	gctgattccg	gtgcagccaa	gcccttacga	catatggg	accgccgacc	7020
	tggtggagct	gggttaagcag	cgcatgagg	tcacggatgg	aaggctacaa	gcgccctttg	7080
	tcgtgtcgcg	ggcgatcaaa	ggcacgcgca	tcggcggtga	ggttgccgag	gcgctggccg	7140
	ggtacgagct	gcccattctt	gagtcccgta	tcacgcagcg	cgtgagctac	ccaggcactg	7200
	ccgccgcgg	cacaaccgtt	cttgaatcag	aaccgcagg	cgacgctgcc	cgcgaggctc	7260
20	aggcgctggc	cgctgaaatt	aaatcaaaac	tcatttgagt	taatgaggta	aagagaaaat	7320
	gagcaaaagc	acaaacacgc	taagtgccgg	ccgtccgagc	gcacgcagca	gcaaggctgc	7380
	aacgttggcc	agcctggcag	acacgccagc	catgaagcgg	gtcaactttc	agttgccggc	7440
	ggaggatcac	accaagctga	agatgtacgc	ggtacgccaa	ggcaagacca	ttaccgagct	7500
	gctatctgaa	tacatcgcg	agctaccaga	gtaaattgagc	aaatgaataa	atgagtagat	7560
25	gaatttttagc	ggctaaagga	ggcggcatgg	aaaatcaaga	acaacaggc	accgacgccg	7620
	tggaatgcc	catgtgtgga	ggaacggg	gttgggcagg	cgtaagcggc	tgggttgtct	7680
	gccggccctg	caatggcact	ggaaccccc	agcccagagg	atcggcgatga	gcggtcgcaa	7740
	accatccggc	ccggtacaaa	tcggcgcg	gctgggtgat	gacctggtgg	agaagttgaa	7800
	ggccgcgcag	gccgcccagc	ggcaacgc	cgaggcgagaa	gacgcccc	tgaatcgtgg	7860
30	caaggggccc	ctgatcgaat	ccgcaaagaa	tcccggcaac	cgccggcagc	cgggtgcgccg	7920
	tcgattagga	agccgccc	gggcgacgag	caaccagatt	ttttcgttcc	gatgctctat	7980
	gacgtgggca	cccgcgatag	tcgcagcatc	atggacgtgg	ccgttttccg	tctgtcgaag	8040
	cgtgaccgac	gagctggcga	ggtgatccgc	tacgagcttc	cagacgggca	cgtagaggtt	8100
	tccgcagggc	cggccggcat	ggccagtgtg	tgggattacg	acctggtact	gatggcggtt	8160
35	tcccatctaa	ccgaatccat	gaaccgatac	cgggaaggga	aggagagaaa	gcccggccgc	8220
	gtgttccgtc	cacacgttgc	ggacgtactc	aagtcttgcc	ggcgagccga	tggcggaag	8280
	cagaaagacg	acctggtaga	aacctgcatt	cggttaaaca	ccacgcacgt	tgccatgcag	8340
	cgtacgaaga	aggccaagaa	cggccgcctg	gtgacggtat	ccgagggtga	agccttgatt	8400
	agccgctaca	agatcgtaaa	gagcgaaacc	ggcgggccgg	agtacatcga	gatcgagcta	8460
40	gctgattgga	tgtaccgcga	gatcacagaa	ggcaagaacc	cggacgtgct	gacggttcac	8520
	cccgattact	ttttgatcga	tcccggcatc	ggccgttttc	tctaccgctt	ggcacgcgc	8580
	gccgcaggca	aggcagaagc	cagatggttg	ttcaagacga	tctacgaacg	cagtggcagc	8640
	gccggagagt	tcaagaagtt	ctgtttcacc	gtgcgcaagc	tgatcgggtc	aaatgacctg	8700
	ccggagtagc	atgtgaagga	ggaggcgggg	caggctggcc	cgatcctagt	catgcgctac	8760
45	cgcaacctga	tcgagggcga	agcatccgcc	ggttccta	gtacggagca	gatgctaggg	8820
	caaattgccc	tagcagggga	aaaaggtcga	aaaggtctct	ttcctgtgga	tagcacgtac	8880
	attgggaacc	caaagccgta	cattgggaac	cggaaaccgt	acattgggaa	cccaaagccg	8940
	tacattggga	accggtcaca	catgtaagt	actgatataa	aagagaaaaa	aggcgatttt	9000
	tccgcctaaa	actctttaaa	acttattaaa	actcttaaaa	cccgcctggc	ctgtgcataa	9060



	ctgtctggcc agcgcacagc cgaagagctg caaaaagcgc ctacccttcg gtcgctgcgc	9120
	tccctacgcc ccgccgcttc gcgtcggcct atcgcggcct atgcgggtgtg aaataccgca	9180
	cagatgcgta aggagaaaat accgcatcag gcgctcttcc gcttcctcgc tcaactgactc	9240
5	gctgcgctcg gtcgttcggc tgcggcgagc ggtatcagct cactcaaagg cggtaatacg	9300
	gttatccaca gaatcagggg ataacgcagg aaagaacatg tgagcaaaag gccagcaaaa	9360
	ggccaggaac cgtaaaaagg ccgcgttgct ggcgtttttc cataggctcc gccccctga	9420
	cgagcatcac aaaaatcgac gctcaagtca gaggtggcga aaccgcagag gactataaag	9480
	ataccaggcg tttccccctg gaagctccct cgtgcgctct cctggtccga ccctgccgct	9540
10	taccggatac ctgtccgcct ttctcccttc gggaaagcgtg gcgctttctc atagctcacg	9600
	ctgtaggtat ctcagttcgg tgtaggtcgt tcgctccaag ctgggctgtg tgcacgaacc	9660
	ccccgttcag cccgaccgct gcgccttctc cggttaactat cgtcttgagt ccaaccgggt	9720
	aagacacgac ttatcgccac tggcagcagc cactggtaac aggattagca gagcgaggta	9780
	tgtaggcggt gctacagagt tcttgaagtg gtggcctaac tacggctaca ctagaaggac	9840
	agtatttggt atctgcgctc tgcgtgaagc agttaccttc ggaaaaagag ttggtagctc	9900
15	ttgatccggc aaacaaacca ccgctggtag cgggtggtttt tttgtttgca agcagcagat	9960
	tacgcgcaga aaaaaaggat ctcaagaaga tcctttgatc ttttctacgg ggtctgacgc	10020
	tcagtggaaac gaaaactcac gttaagggat tttggtcatg catgatatat ctcccaattt	10080
	gtgtagggct tattatgcac gcttaaaaat aataaaagca gacttgacct gatagtttg	10140
20	ctgtgagcaa ttatgtgctt agtgcacta acgcttgagt taagccgcgc cgcgaagcgg	10200
	cgtcggcttg aacgaatttc tagctagaca ttatttgccg actaccttg tgatctcgcc	10260
	tttcacgtag tggacaaatt cttccaactg atctgcgcgc gaggccaaagc gatcttcttc	10320
	ttgtccaaga taagcctgtc tagcttcaag tatgacgggc tgatactggg ccggcaggcg	10380
	ctccattgcc cagtcggcag cgacatcctt cggcgcgatt ttgccggtta ctgcgctgta	10440
	ccaaatgcgg gacaacgtaa gcactacatt tcgctcatcg ccagcccagt cgggcggcga	10500
25	gttccatagc gttaagggtt catttagcgc ctcaaataga tcctgttcag gaaccggatc	10560
	aaagagttcc tccgccgctg gacctaccaa ggcaacgcta tgttctcttg cttttgtcag	10620
	caagatagcc agatcaatgt cgatcgtggc tggctcgaag atacctgcaa gaatgtcatt	10680
	gcgctgccat tctccaaatt gcagttcgcg cttagctgga taacgccacg gaatgatgtc	10740
30	gtcgtgcaca acaatggtga cttctacagc gcggagaatc tcgctctctc cagggggaagc	10800
	cgaagtttcc aaaaggctgt tgatcaaagc tcgccgcggt gtttcatcaa gccttacggg	10860
	caccgtaacc agcaaatcaa tatcactgtg tggcttcagg ccgccatcca ctgcggagcc	10920
	gtacaaatgt acggccagca acgtcgggtc gagatggcgc tcgatgacgc caactacctc	10980
	tgatagttga gtcgatactt cggcgatcac cgcttcccc atgatgttta actttgtttt	11040
	agggcgactg ccctgctgcg taacatcgtt gctgctccat aacatcaaac atcgaccac	11100
35	ggcgtaacgc gcttgctgct tggatgccc aggcatagac tgtaccccaa aaaaacagtc	11160
	ataacaagcc atgaaaaccg ccactgcgtt ccatg	11195
	<210> 3	
40	<211> 12415	
	<212> DNA	
	<213> synthetic: expression vector - binary vector pSUNubiGUS	
	<400> 3	
45	aatattcaaa caaacacata cagcgcgact tatcatggac atacaaatgg acgaacggat	60
	aaaccttttc acgccctttt aaatatccga ttattctaat aaacgctctt ttctcttagg	120
	tttaccgcc aatatatcct gtcaaacact gatagtttaa actgaaggcg ggaaacgaca	180
	atcagatcta gtaggaaaca gctatgacca tgattacgcc aagcttgcac gcctgcagg	240
	cgaaattcga gtcgggtacc cggtcactgg attttggtt taggaattag aaattttatt	300

	gatagaagta	ttttacaaat	acaaatacat	actaaggggt	tcttatatgc	tcaacacatg	360
	agcgaaaccc	tataagaacc	ctaattccct	tatctgggaa	ctactcacac	attattctgg	420
	agaaaaatag	agagagatag	atthttagag	agagactgg	gattttttgc	cgggtaccg	480
	agctcggtag	caattcccga	ggctgtagcc	gacgatggg	cgccaggaga	gttggtgatt	540
5	cattgtttgc	ctccctgctg	cggtttttca	ccgaagttca	tgccagtcca	gcgtttttgc	600
	agcagaaaag	ccgccgactt	cggtttgccg	tcgcgagtga	agatcccttt	cttggtaccg	660
	ccaacgcgca	atatgccttg	cgaggtcgca	aaatcggcga	aattccatac	ctgttcaccg	720
	acgacggcgc	tgacgcgac	aaagacgcg	tgatacatat	ccagccatgc	acactgatac	780
	tcttcactcc	acatgtcgg	gtacattgag	tgacgcccgg	ctaacgtatc	cacgccgtat	840
10	tcgggtgatga	taatcggctg	atgcagtttc	tcttgccagg	ccagaagtcc	tttttccagt	900
	accttctctg	ccgtttccaa	atcgccgctt	tgacataacc	atccgtaata	acgggttcagg	960
	cacagcacat	caaagagatc	gctgatggta	tcgggtgtgag	cgtcgcagaa	cattacattg	1020
	acgcaggtga	tcggacgcgt	cgggtcgagt	ttacgcggtg	cttcgccag	tgccgaaata	1080
	ttcccgtgca	cttgccgacg	ggatccgg	tcgttgccaa	tactccacat	caccacgctt	1140
15	gggtgggtttt	tgacacgcgc	tatcagctct	ttaatcgctt	gtaagtgcgc	ttgctgagtt	1200
	tccccgttga	ctgcctcttc	gctgtacagt	tctttcggct	tggtgcccgc	ttcgaaacca	1260
	atgcctaaag	agaggttaaa	gccgacagca	gcagtttcat	caatcaccac	gatgccatgt	1320
	tcactctgcc	agtcgagcat	ctcttcagcg	taagggtaat	gcgaggtacg	gtaggagttg	1380
	gccccaatcc	agtcatttaa	tgctgggtcg	tgacccatca	gcacgttata	gaatcccttg	1440
20	ccacgtaagt	ccgcattctt	atgacgacca	aagccagtaa	agtagaacgg	tttggtggtta	1500
	atcaggaact	gttcgccctt	cactgccact	gaccggatgc	cgacgcgaag	cgggtagata	1560
	tcacactctg	tctggctttt	ggctgtgacg	cacagttcat	agagataacc	ttcaccgggt	1620
	tgccagaggt	gcggattcac	cacttgcaaa	gtcccgcctg	tgcttctgtc	agttgcaacc	1680
	acctgttgat	ccgcattcac	cagttcaaac	ctgacatcac	cattggccac	cacctgccag	1740
25	tcaacagacg	cgtgggttaca	gtcttgccgc	acatgcgtca	ccacgggtgat	atcgctccacc	1800
	caggtgttcg	gcgtgggtgta	gagcattacg	ctgcgatgga	ttccggcata	gttaaagaaa	1860
	tcattggaagt	aagactgctt	tttcttgccg	ttttcgtcgg	taatcaccat	tccggcgggg	1920
	atagtctgcc	agttcagttc	gttggttcaca	caaacgggtga	tacctgcaca	tcaacaaatt	1980
	ttggtcatat	attagaaaag	ttataaaatta	aaatatacac	acttataaac	tacagaaaag	2040
30	caattgctat	atactacatt	ctttttatttt	gaaaaaaata	tttgaaatat	tattattacta	2100
	ctaattaatg	ataattatta	tatatatatc	aaaggtagaa	gcagaaactt	acgtacactt	2160
	ttcccggcaa	taacatacgg	cgtgacatcg	gcttcaaatg	gcgtatagcc	gccctgatgc	2220
	tccatcactt	cctgattatt	gaccacactt	ttgccgtaat	gagtgaccgc	atcgaaacgc	2280
	agcacgatac	gctggcctgc	ccaacctttc	ggtataaaaga	cttcgcgctg	ataccagacg	2340
35	ttgcccgcat	aattacgaat	atctgcatcg	gcgaactgat	cgttaaaact	gcctggcaca	2400
	gcaattgccc	ggctttcttg	taacgcgctt	tcccaccaac	gctgaccaat	tccacagttt	2460
	tcgcgatcca	gactgaatgc	ccacaggccg	tcgagttttt	tgatttcacg	ggttgggggt	2520
	tctacaggac	gtaccatgg	cgatcgactc	tagactagtg	gatccgatat	cgcccgggct	2580
	gcagaagtaa	caccaaaaca	cagggtgagc	atcgacaaaa	gaaacagtac	caagcaaata	2640
40	aatagcgtat	gaaggcagg	ctaaaaaaat	ccacatatag	ctgctgcata	tgccatcatc	2700
	caagtatatc	aagatcaaaa	taattataaa	acatacttgt	ttattataat	agataggtac	2760
	tcaaggttag	agcatatgaa	tagatgctgc	atatgccatc	atgtatatgc	atcagtaaaa	2820
	cccacatcaa	catgtatacc	tatcctagat	cgatatttcc	atccatctta	aactcgtaac	2880
	tatgaagatg	tatgacacac	acatacagtt	ccaaaattaa	taaatacacc	aggtagtttg	2940
45	aaacagtatt	ctactccgat	ctagaacgaa	tgaacgaccg	cccaaccaca	ccacatcatc	3000
	acaaccaagc	gaacaaaaag	catctctgta	tatgcatcag	taaaaccgcg	atcaacatgt	3060
	atacctatcc	tagatcgata	tttccatcca	tcatcttcaa	ttcgtaacta	tgaatatgta	3120
	tgccacacac	atacagatcc	aaaattaata	aatccaccag	gtagtttgaa	acagaattct	3180
	actccgatct	agaacgaccg	cccaaccaga	ccacatcatc	acaaccaaga	caaaaaaaag	3240

	catgaaaaga	tgacccgaca	aacaagtgca	cggcatatat	tgaaataaag	gaaaagggca	3300
	aaccaaacc	tatgcaacga	aacaaaaaaa	atcatgaaat	cgatcccgtc	tgcggaacgg	3360
	ctagagccat	cccaggattc	cccaaagaga	aacactggca	agtttagcaat	cagaacgtgt	3420
	ctgacgtaca	ggtcgcatcc	gtgtacgaac	gctagcagca	cggatctaac	acaaacacgg	3480
5	atctaacaca	aacatgaaca	gaagtagaac	taccgggccc	taaccatgga	ccggaacgcc	3540
	gatctagaga	aggtagagag	gggggggggg	ggaggacgag	cggcgtaact	tgaagcggag	3600
	gtgccgacgg	gtggatttgg	gggagatctg	gttgtgtgtg	tgtgcgctcc	gaacaacacg	3660
	aggttgggga	aagaggggtg	ggaggggggtg	tctattttatt	acggcggggcg	aggaagggaa	3720
	agcgaaggag	cgggtgggaaa	ggaatcccc	gtagctgccg	tgccgtgaga	ggaggaggag	3780
10	gccgcctgcc	gtgccggctc	acgtctgccg	ctccgccacg	caatttctgg	atgccgacag	3840
	cggagcaagt	ccaacgggtg	agcggaaactc	tcgagagggg	tccagaggca	gcgacagaga	3900
	tgccgtgccg	tctgcttcgc	ttggcccagc	gcgacgctgc	tggttcgctg	gttgggtgtcc	3960
	gttagactcg	tcgacggcgt	ttaacaggct	ggcattatct	actcgaaaca	agaaaaatgt	4020
	ttccttagtt	tttttaattt	cttaaagggt	atttgtttta	tttttagtca	ctttattttta	4080
15	ttctatttta	tatctaaatt	attaaataaa	aaaactaaaa	tagagtttta	gttttcttaa	4140
	tttagaggct	aaaatagaat	aaaatagatg	tactaaaaaa	attagtctat	aaaaaccatt	4200
	aaccctaacc	cctaaatgga	tgtactaata	aaatggatga	agtattatat	aggtgaagct	4260
	atttgcaaaa	aaaagaggaga	acacatgcac	actaaaaaga	taaaactgta	gagtcctgtt	4320
	gtcaaaaatac	tcaattgtcc	tttagaccat	gtctaactgt	tcatttatat	gattctctaa	4380
20	aacactgata	ttattgtagt	actatagatt	atattattcg	tagagtaaag	tttaaatata	4440
	tgtataaaga	tagataaaact	gcacttcaaa	caagtgtgac	aaaaaaaaata	tgtggtaatt	4500
	ttttataact	tagacatgca	atgctcatta	tctctagaga	ggggcacgac	cgggtcacgc	4560
	tgcaactgcag	gaattcgatg	ggctcgaggt	accgagctcg	gtacccactg	gattttgggt	4620
	tttaggaatta	gaaattttat	tgatagaagt	attttacaaa	tacaaatata	tactaagggt	4680
25	ttcttatatg	ctcaacacat	gagcgaaacc	ctataagaac	cctaattccc	ttatctggga	4740
	actactcaca	cattattata	gagagagata	gatttgtaga	gagagactgg	tgatttcagc	4800
	ggcatgcctg	caggtcgact	cagatctggg	taactggcct	aactggcctt	ggaggagctg	4860
	gcaactcaaa	atccctttgc	caaaaaccaa	catcatgcca	tccaccatgc	ttgtatccag	4920
	ctgcgcgcaa	tgtaccccg	gctgtgtatc	ccaaagcctc	atgcaacctc	acagatggat	4980
30	cgtttggaag	gcctataaca	gcaaccacag	acttaaaacc	ttgcgcctcc	atagacttaa	5040
	gcaaatgtgt	gtacaatgtg	gatcctaggc	ccaacctttg	atgcctatgt	gacacgtaaa	5100
	cagtactctc	aactgtccaa	tcgtaagcgt	tcttagcctt	ccagggccca	gcgtaagcaa	5160
	taccagccac	aacaccctca	acctcagcaa	ccaaccaagg	gtatctatct	tgcaacctct	5220
	ctagatcatc	aatccactct	tgtggtgttt	gtggctctgt	cctaaagtcc	actgtagacg	5280
35	tctcaatgta	atggttaacg	atatcaca	ccgcggccat	atcagctgct	gtagctggcc	5340
	taatctcaac	tggtctcctc	tccggagaca	tgtcgactct	agaggatccc	ccgggctgca	5400
	gaagtaacac	caaacaacag	ggtgagcatc	gacaaaagaa	acagtaccaa	gcaataaat	5460
	agcgtatgaa	ggcagggcta	aaaaaatcca	catatagctg	ctgcatatgc	catcatccaa	5520
	gtatatcaag	atcaaaataa	ttataaaaaca	tacttgttta	ttataataga	taggtactca	5580
40	aggttagagc	atatgaatag	atgctgcata	tgccatcatg	tatatgcac	agtaaaaccc	5640
	acatcaacat	gtatacctat	cctagatcga	tatttccatc	catcttaaac	tcgtaactat	5700
	gaagatgtat	gacacacaca	tacagttcca	aaattaataa	atacaccagg	tagtttgaaa	5760
	cagtattcta	ctccgatcta	gaacgaatga	acgaccgccc	aaccacacca	catcatcaca	5820
	accaagcgaa	caaaaagcat	ctctgtatat	gcatcagtaa	aaccgcgcatc	aacatgtata	5880
45	cctatcctag	atcgatatatt	ccatccatca	tcttcaattc	gtaactatga	atatgtatgg	5940
	cacacacata	cagatccaaa	attaataaat	ccaccaggta	gtttgaaaca	gaattctact	6000
	ccgatctaga	acgaccgccc	aaccagacca	catcatcaca	accaagacaa	aaaaaagcat	6060
	gaaaagatga	cccgacaaac	aagtgcacgg	catatatattga	aataaaggaa	aagggcaaac	6120
	caaaccctat	gcaacgaaac	aaaaaaaaatc	atgaaatcga	tcccgctctgc	ggaacggcta	6180

	gagccatccc	aggattcccc	aaagagaaac	actggcaagt	tagcaatcag	aacgtgtctg	6240
	acgtacaggt	cgcacccgtg	tacgaacgct	agcagcacgg	atctaacaca	aacacggatc	6300
	taacacaaaac	atgaacagaa	gtagaactac	cgggccctaa	ccatggaccg	gaacgccgat	6360
	ctagagaagg	tagagagggg	ggggggggga	ggacgagcgg	cgtaccttga	agcggagggtg	6420
5	ccgacgggtg	gatttggggg	agatctggtt	gtgtgtgtgt	gcgctccgaa	caacacgagg	6480
	ttggggaaaag	aggggtgtgga	gggggtgtct	atattattacg	gcgggagagg	aagggaaagc	6540
	gaaggagcgg	tgggaaaagga	atcccccgta	gctgccgtgc	cgtgagagga	ggaggaggcc	6600
	gcctgccgtg	ccggctcacg	tctgccgctc	cggcacgcaa	tttctggatg	ccgacagcgg	6660
	agcaagtcca	acgggtggagc	ggaactctcg	agaggggtcc	agaggcagcg	acagagatgc	6720
10	cgtgccgtct	gcttcgcttg	gcccgcgcgc	acgctgctgg	ttcgctgggt	ggtgtccgtt	6780
	agactcgtcg	acggcggtta	acaggctggc	attatctact	cgaacaaga	aaaatgtttc	6840
	cttagttttt	ttaattttctt	aaagggtatt	tgtttaattt	ttagtcactt	tattttattc	6900
	tattttatat	ctaaattatt	aaataaaaaa	actaaaaatag	agtttttagt	ttcttaattt	6960
	agaggctaaa	atagaataaa	atagatgtac	taaaaaaatt	agtctataaa	aaccattaac	7020
15	cctaaaccct	aaatggatgt	actaataaaa	tggatgaagt	attatatagg	tgaagctatt	7080
	tgcaaaaaaa	aaggagaaca	catgcacact	aaaaagataa	aactgtagag	tcctgttgtc	7140
	aaaataactca	attgtccttt	agaccatgtc	taactgttca	tttatatgat	tctctaaaac	7200
	actgatatta	ttgtagtact	atagattata	ttattcgtag	agtaaaagtt	aaatatatgt	7260
	ataaagatag	ataaactgca	cttcaaacaa	gtgtgacaaa	aaaaaatatgt	ggtaattttt	7320
20	tataacttag	acatgcaatg	ctcattatct	ctagagaggg	gcacgaccgg	gtcacgctgc	7380
	actgcaggaa	ttcgatatca	agctggctga	gtggctcctt	caacgttgcg	gttctgtcag	7440
	ttccaaacgt	aaaacggctt	gtcccgcgtc	atcggcgggg	gtcataacgt	gactccctta	7500
	attctccgct	catgatcaga	ttgtcggttc	ccgccttcag	tttaaaactat	cagtgtttga	7560
	caggatcact	gcttggtaat	aattgtcatt	agattgtttt	tatgcataga	tgcactcgaa	7620
25	atcagccaat	tttagacaag	tatcaaaccg	atgttaattc	agtacattaa	agacgtccgc	7680
	aatgtgttat	taagttgtct	aagcgtcaat	ttgtttacac	cacaatatat	cctgccacca	7740
	gccagccaac	agctccccga	ccggcagctc	ggcacaaaat	caccacgcgt	ctaaaaaggt	7800
	gatgtgtatt	tgagtaaaac	agcttgctgc	atgcggctgc	tgcttatatg	atgcgatgag	7860
	taaataaaca	aatacgcaag	gggaacgcat	gaagggtatc	gctgtactta	accagaaagg	7920
30	cgggtcaggc	aagacgacca	tgcgaaccca	tctagcccgc	gccctgcaac	tcgccggggc	7980
	cgatgttctg	ttagtogatt	ccgatcccca	gggcagtgcc	cgcgattggg	cggccgtgcg	8040
	ggaagatcaa	ccgctaaccg	ttgtcggcat	cgaccgcccg	acgattgacc	gcgacgtgaa	8100
	ggccatcggc	cggcgcgact	tcgtagtgat	cgacggagcg	ccccaggcgg	cggacttggc	8160
	tgtgtccgcg	atcaaggcag	ccgacttcgt	gctgattccg	gtgcagccaa	gcccttacga	8220
35	catatggggc	accgcccagc	tgggtggagct	ggttaagcag	cgcattgagg	tcacggatgg	8280
	aaggctacaa	gcggcccttg	tcgtgtcgcg	ggcgatcaaa	ggcacgcgca	tcggcgggtga	8340
	ggttgccgag	gcgctggccg	ggtacgagct	gcccattctt	gagtcccgtg	tcacgcagcg	8400
	cgtgagctac	ccaggcactg	ccgcccgcgg	cacaaccgtt	cttgaatcag	aacccgaggg	8460
	cgacgctgcc	cgcgagggtc	aggcgctggc	cgctgaaatt	aaatcaaaac	tcatttgagt	8520
40	taatgaggta	aagagaaaaat	gagcaaaaagc	acaaacacgc	taagtgccgg	ccgtccgagc	8580
	gcacgcagca	gcaaggctgc	aacggttgcc	agcctggcag	acacgccagc	catgaagcgg	8640
	gtcaactttc	agttgccggc	ggaggatcac	accaagctga	agatgtacgc	ggtacgccaa	8700
	ggcaagacca	ttaccgagct	gctatctgaa	tacatcgcg	agctaccaga	gtaaatgagc	8760
	aaatgaataa	atgagtagat	gaatttttagc	ggctaaagga	ggcggcatgg	aaaatcaaga	8820
45	acaaccaggc	accgacgccg	tgggaatgcc	catgtgtgga	ggaacgggcg	gttggccagg	8880
	cgtaagcggc	tgggttgtct	gccggccctg	caatggcact	ggaaccccc	agcccagagga	8940
	atcggcgtga	gcggctcgca	accatccggc	ccggtacaaa	tcggcgcggc	gctgggtgat	9000
	gacctggtgg	agaagttgaa	ggccgcgcag	gccgcccagc	ggcaacgcat	cgaggcagaa	9060
	gacgccccgg	tgaatcgtgg	caagggggccg	ctgatcgaat	ccgcaaagaa	tcccggcaac	9120

	cgccggcagc	cggtgcgccc	tcgattagga	agccgccc	gggagcag	caaccagatt	9180
	ttttcgttcc	gatgctctat	gacgtgggca	cccgcgatag	tcgcagcatc	atggacgtgg	9240
	ccgtttttccg	tctgtcgaag	cgtgaccgac	gagctggcga	ggtgatccgc	tacgagcttc	9300
	cagacgggca	cgtagagggt	tccgcagggc	cggccggcat	ggccagtgtg	tgggattacg	9360
5	acctgggtact	gatggcggtt	tcccattctaa	ccgaatccat	gaaccgatac	cgggaaggga	9420
	aggagagacaa	gcccggccgc	gtgttccgtc	cacacgttgc	ggacgtactc	aagttctgcc	9480
	ggcgagccga	tggcggaag	cagaaagacg	acctggtaga	aacctgcatt	cggttaaaca	9540
	ccacgcacgt	tgccatgcag	cgtacgaaga	aggccaagaa	cggccgcctg	gtgacgggat	9600
	ccgaggggtga	agccttgatt	agccgctaca	agatcgtaaa	gagcgaaaacc	gggaggcccg	9660
10	agtacatcga	gatcgagcta	gctgattgga	tgtaccgcga	gatcacagaa	ggcaagaacc	9720
	cggacgtgct	gacgggtcac	cccgattact	ttttgatcga	tcccggcatc	ggcgttttc	9780
	tctaccgcct	ggcagccgc	gccgcaggca	aggcagaagc	cagatgggtg	ttcaagacga	9840
	tctacgaacg	cagtggcagc	gccggagagt	tcaagaagtt	ctgtttcacc	gtgcgcaagc	9900
	tgatcgggtc	aaatgacctg	ccggagtacg	atgtgaagga	ggaggcgggg	caggctggcc	9960
15	cgatccctagt	catgcgctac	cgcaacctga	tcgagggcga	agcatccgcc	ggttcctaata	10020
	gtacggagca	gatgctaggg	caaattgccc	tagcagggga	aaaaggctcg	aaaggctctct	10080
	ttcctgtgga	tagcacgtac	attgggaacc	caaagccgta	cattgggaac	cggaaaccgt	10140
	acattgggaa	cccaagccg	tacattggga	accggtcaca	catgtaagt	actgatataa	10200
	aagagaaaaa	aggcgatttt	tccgcctaaa	actctttaaa	acttattaaa	actcttaaaa	10260
20	cccgcctggc	ctgtgcataa	ctgtctggcc	agcgcacagc	cgaagagctg	caaaaagcgc	10320
	ctacccttcg	gtcgtcgcgc	tccctacgcc	ccgcgccttc	gcgtcggcct	atcgcggcct	10380
	atgcgggtgtg	aaataccgca	cagatgcgta	aggagaaaat	accgcacatc	gcgtctcttc	10440
	gcttccctgc	tactgaactc	gctgcgctcg	gtcgttcggc	tgccggcagc	ggtatcagct	10500
	cactcaaagg	cggtaatagc	gttatccaca	gaatcagggg	ataacgcagg	aaagaacatg	10560
25	tgagcaaaag	gccagcaaaa	ggccaggaac	cgtaaaaagg	ccgcgttgct	ggcgtttttc	10620
	cataggctcc	gccccctga	cgagcatcac	aaaaatcgac	gctcaagtca	gaggtggcga	10680
	aaccgcagag	gactataaag	ataccaggcg	tttccccctg	gaagctccct	cgtgcgctct	10740
	cctgtttccga	ccctgccgct	taccggatac	ctgtccgcct	ttctcccttc	gggaagcgtg	10800
	gcgctttctc	atagctcagc	ctgtaggtat	ctcagttcgg	tgtaggtcgt	tcgctccaag	10860
30	ctgggctgtg	tgcacgaacc	cccgttcag	ccgaccgct	gcgccttatc	cggtaactat	10920
	cgtcttgagt	ccaaccgggt	aagacacgac	ttatcgccac	tggcagcagc	cactggtaac	10980
	aggattagca	gagcgaggta	tgtaggcggt	gctacagagt	tcttgaagt	gtggcctaac	11040
	tacggctaca	ctagaaggac	agtatttggt	atctgcgctc	tgctgaagcc	agttaccttc	11100
	ggaaaaagag	ttggtagctc	ttgatccggc	aaacaaacca	ccgctggtag	cgggtggtttt	11160
35	tttggttgca	agcagcagat	tacgcgcaga	aaaaaaggat	ctcaagaaga	tcctttgatc	11220
	ttttctacgg	ggtctgacgc	tcagtggaa	gaaaactcac	gttaagggat	tttggtcatg	11280
	catgatata	ctccaattt	gtgtagggt	tattatgcac	gcttaaaaat	aataaaagca	11340
	gacttgacct	gatagtttgg	ctgtgagcaa	ttatgtgctt	agtgcattca	acgcttgagt	11400
	taagccgcgc	cgcgaagcgg	cgtcggcttg	aacgaatttc	tagctagaca	ttatttgccg	11460
40	actaccttg	tgatctcgcc	tttcacgtag	tggacaaaat	cttccaactg	atctgcgcgc	11520
	gaggccaagc	gatcttcttc	ttgtccaaga	taagcctgtc	tagcttcaag	tatgacgggc	11580
	tgatactggg	ccggcaggcg	ctccattgcc	cagtcggcag	cgacatcctt	cggcgcgatt	11640
	ttgccgggta	ctgcgctgta	ccaaatgcgg	gacaacgtaa	gcactacatt	tcgctcatcg	11700
	ccagcccagt	cgggcggcga	gttccatagc	gttaagggtt	catttagcgc	ctcaaataga	11760
45	tcctgttcag	gaaccggatc	aaagagttcc	tccgcgcgtg	gacctaccaa	ggcaacgcta	11820
	tgttctcttg	cttttgtcag	caagatagcc	agatcaatgt	cgatcgtggc	tggctcgaag	11880
	atacctgcaa	gaatgtcatt	gcgctgccat	tctccaaatt	gcagttcgcg	cttagctgga	11940
	taacgccacg	gaatgatgtc	gtcgtgcaca	acaatgggtga	cttctacagc	gcggagaatc	12000
	tcgctctctc	caggggaagc	cgaagtcttc	aaaaggctcg	tgatcaaagc	tcgccgcgtt	12060

gtttcatcaa gccttacggt caccgtaacc agcaaataca tatcactgtg tggcttcagg 12120  
ccgccatcca ctgcggagcc gtacaaatgt acggccagca acgtcgggtc gagatggcgc 12180  
tcgatgacgc caactacctc tgatagttga gtcgatactt cggcgatcac cgcttcccc 12240  
atgatgttta actttgtttt agggcgactg ccctgctgcg taacatcggt gctgctccat 12300  
5 aacatcaaac atcgaccac gccgtaacgc gcttgctgct tggatgcccg aggcataagac 12360  
tgtaccccaa aaaaacagtc ataacaagcc atgaaaaccg ccactgcggt ccatg 12415

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2004/007567

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N15/82 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, EMBASE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 97/48814 A (MONSANTO CO) 24 December 1997 (1997-12-24) cited in the application page 26 - page 35	1-12
Y	KUMLEHN JOCHEN ET AL: "Differentiation of isolated wheat zygotes into embryos and normal plants" PLANTA (BERLIN), vol. 205, no. 3, July 1998 (1998-07), pages 327-333, XP002263491 ISSN: 0032-0935 cited in the application the whole document	1-12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

27 October 2004

Date of mailing of the international search report

17/11/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Bucka, A

## INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/EP2004/007567

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	HOLM P B ET AL: "TRANSFORMATION OF BARLEY BY MICROINJECTION INTO ISOLATED ZYGOTE PROTOPLASTS" TRANSGENIC RESEARCH, LONDON, GB, vol. 9, February 2000 (2000-02), pages 21-32, XP000945396 ISSN: 0962-8819 the whole document	1-12
A	US 6 300 543 B1 (MCGANN LOCKSLEY E ET AL) 9 October 2001 (2001-10-09) examples 1-3,7	1-12
A	KRANZ E ET AL: "ANGIOSPERM FERTILISATION, EMBRYO AND ENDOSPERM DEVELOPMENT IN VITRO" PLANT SCIENCE, LIMERICK, IE, vol. 142, 1999, pages 183-197, XP002935311 ISSN: 0168-9452 the whole document	1-12
A	WO 01/73084 A (INST PFLANZENGENETIK & KULTUR ; POPELKA JUAN CARLOS (DE); ALTPETER) 4 October 2001 (2001-10-04) page 15 - page 16	1-12
A	RASCO-GAUNT SONRIZA ET AL: "Procedures allowing the transformation of a range of European elite wheat (Triticum aestivum L.) varieties via particle bombardment" JOURNAL OF EXPERIMENTAL BOTANY, vol. 52, no. 357, April 2001 (2001-04), pages 865-874, XP002263635 ISSN: 0022-0957 the whole document	1-12
A	FRAME BRONWYN R ET AL: "Agrobacterium tumefaciens-mediated transformation of maize embryos using a standard binary vector system" PLANT PHYSIOLOGY (ROCKVILLE), vol. 129, no. 1, May 2002 (2002-05), pages 13-22, XP002263971 ISSN: 0032-0889 the whole document	1-12



## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP2004/007567

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9748814	A	24-12-1997	AU 738153 B2	13-09-2001
			AU 3402897 A	07-01-1998
			CA 2230216 A1	24-12-1997
			CN 1208437 A	17-02-1999
			CZ 9800867 A3	17-03-1999
			EP 0856060 A2	05-08-1998
			HU 9902123 A2	28-10-1999
			TR 9800294 T1	21-10-1999
			WO 9748814 A2	24-12-1997
			US 2003024014 A1	30-01-2003
US 6300543	B1	09-10-2001	AU 3495297 A	02-02-1998
			AU 731406 B2	29-03-2001
			AU 3718697 A	02-02-1998
			CA 2259983 A1	15-01-1998
			EP 0914455 A1	12-05-1999
			WO 9801575 A1	15-01-1998
			WO 9801576 A1	15-01-1998
			ZA 9706074 A	14-04-1999
WO 0173084	A	04-10-2001	DE 10015458 A1	11-10-2001
			AU 7382101 A	08-10-2001
			WO 0173084 A2	04-10-2001
			DE 10115761 A1	20-12-2001

**THIS PAGE BLANK (USPTO)**